

Customer Number: 000959

**DIVISIONAL-CONTINUATION APPLICATION TRANSMITTAL FORM****UNDER RULE 1.53(b) (former Rule 1.60)**

DOCKET NUMBER	ANTICIPATED CLASSIFICATION OF THIS APPLICATION:		PRIOR APPLICATION SERIAL NUMBER: 08/636,602	PRIOR APPLICATION FILING DATE: APRIL 23, 1996
HUI-021CN	CLASS:	SUBCLASS:	EXAMINER: S. PRIEBE	ART UNIT: 1632

ASSISTANT COMMISSIONER FOR PATENTS  
BOX PATENT APPLICATION  
WASHINGTON, DC 20231

**CERTIFICATION UNDER 37 CFR 1.10**

Date of Deposit: February 17, 1999

Mailing Label Number: EL094334226US

I hereby certify that this 37 CFR 1.53(b) request and the documents referred to as attached therein are being deposited with the United States Postal Service on the date indicated above in an envelope as "Express Mail Post Office to Addressee" service under 37 CFR 1.10 and addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.



Name of Person Mailing Paper

Signature of Person Mailing Paper

Dear Sir:

This is a request for filing a ☒ continuation ☐ divisional application under 37 CFR 1.53(b), of pending prior application serial no. 08/636,602 filed on April 23, 1996, of Laurie H. Glimcher and I-Cheng Ho entitled Methods and Compositions For Regulating T Cell Subsets By Modulating Transcription Factor Activity.

- ☒ Enclosed is a copy of the latest inventor signed application, including the oath or declaration as originally filed. The copy of the enclosed papers is as follows:
  - ☒ 51 page(s) of specification
  - ☒ 4 page(s) of claims
  - ☒ 1 page(s) of abstract
  - ☒ 12 sheet(s) of drawings
  - ☒ 5 page(s) of declaration and power of attorney.
- ☒ A verified statement to establish small entity status under 37 CFR 1.9 and 1.27, a copy of which is enclosed, was filed in the prior application and such status is still proper and desired (37 CFR 1.28(a)).
- ☒ The filing fee is calculated below:

	NUMBER OF CLAIMS FILED			NUMBER EXTRA
TOTAL	* 21	MINUS	** 20	= 1
INDEP.	* 2	MINUS	*** 3	= 0

☐ MULTIPLE DEPENDENT CLAIMS

## SMALL ENTITY

RATE	FEE
x 9 =	\$ 9.00
x 39 =	\$0.00
+130 =	\$ 0.00
BASIC FEE	\$380.00
TOTAL	\$389.00

OTHER THAN A  
SMALL ENTITY

RATE	FEE
x 18 =	\$ 0.00
x 78 =	\$ 0.00
+ 260 =	\$ 0.00
BASIC FEE	\$ 0.00
TOTAL	\$0.00

- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment, to Deposit Account No. 12-0080. A duplicate copy of this sheet is enclosed.


5. ☒ A check in the amount of \$389.00 is enclosed for payment of the filing fee.
6. ☒ Cancel in this application original claims 1-34 of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)
7. ☒ A preliminary amendment is enclosed. (Claims added by this amendment have been properly numbered consecutively beginning with the number next following the highest numbered original claims in the prior application.)
8. ☒ The prior application is assigned of record to President and Fellows of Harvard College.
9. ☒ Also enclosed is an Associate Power of Attorney to Megan E. Williams, Ph.D..
10. ☒ The power of attorney in the prior application is to Lahive & Cockfield, LLP.
- a. ☒ The power appears in the original papers in the prior application.
- b. ☐ Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.
- c. ☐ A new power has been executed and is attached.
11. ☒ Address all future communications (May only be completed by applicant, or attorney or agent of record) to Giulio A. DeConti, Jr. at **Customer Number: 000959** whose address is:

Lahive & Cockfield, LLP  
28 State Street  
Boston, Massachusetts 02109

12. ☒ Any requests for extensions of time necessary in a parent application for establishing copendency between this application and a parent application are hereby requested and the Commissioner is authorized to charge any fee associated with such an extension to Deposit Account No. 12-0080.
13. ☒ Pursuant to 37 CFR 1.821(e), the computer readable form of the sequence listing for this new application is to be identical with the computer readable form of application serial no. 08/636,602. Please use the computer readable form of application serial no. 08/636,602 in lieu of filing a duplicate computer readable form in this application. Pursuant to 37 CFR 1.821(f), the content of the paper copy of the sequence listing for this new application and the computer readable form of application serial no. 08/636,602 are the same.

February 12, 1999  
Date

LAHIVE & COCKFIELD, LLP  
28 State Street  
Boston, Massachusetts 02109  
Tel. (617) 227-7400

  
Megan E. Williams, Ph.D.  
Reg. No. 43,270  
☐ inventor(s) ☐ filed under §1.34(a)  
☐ assignee of complete interest  
☒ attorney or agent of record

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the application of: Laurie H. Glimcher et al.

Serial No.: Continuation of 08/636,602

Filed:

For: *METHODS AND COMPOSITIONS FOR  
REGULATING T CELL SUBSETS BY  
MODULATING TRANSCRIPTION FACTOR  
ACTIVITY*

Attorney Docket No.: HUI-021CN

Group Art Unit:

Examiner:

Assistant Commissioner for Patents  
Washington, D.C. 20231

**PRELIMINARY AMENDMENT**

Dear Sir:

Prior to examination, please amend the above-referenced patent application as follows:

In the Specification:

After the title please insert:

-- Cross Reference to Related Applications

This application is a continuation application of U.S.S.N. 08/636,602, filed on April 23, 1996. The contents of that application are specifically incorporated herein by reference.--

In the claims:

Please cancel claims 1-34 without prejudice. Please add the following new claims:

09249756-0199  
662120-95284260

35. **(New)** A method for identifying a compound that modulates an immune response, comprising

providing an indicator composition comprising a maf family protein and a target DNA to which said maf family protein bind, said indicator composition being an indicator cell or an acellular preparation;

contacting the indicator composition with each member of a library of test compounds;

selecting from the library of test compounds a compound of interest that modulates the activity of the maf family protein; and

determining the effect of the compound of interest on an immune response to thereby identify a compound that modulates an immune response.

36. **(New)** The method of claim 35, wherein the maf family protein is c-Maf.

37. **(New)** The method of claim 35, wherein the effect of the compound of interest on an immune response is determined by determining the effect of the compound on expression of a Th2-associated cytokine gene.

38. **(New)** The method of claim 37, wherein the Th2-associated cytokine gene is an interleukin-4 gene.

39. **(New)** The method of claim 35, wherein the effect of the compound of interest on an immune response is determined by determining the effect of the compound on development of T helper type 1 (Th1) or T helper type 2 (Th2) cells.

40. **(New)** The method of claim 35, wherein the maf family protein is selected from the group consisting of v-maf, mafB, Nrl, mafK, mafF, mafG and p18.

41. **(New)** The method of claim 35, wherein the target DNA comprises a regulatory sequence of a Th2-associated cytokine gene.

42. **(New)** The method of claim 41, wherein the Th2-associated cytokine gene is an interleukin-4 gene.

43. **(New)** The method of claim 35, wherein the indicator composition is an indicator cell.

09443720 "95284260

44. **(New)** The method of claim 43, wherein the indicator cell is a lymphoid cell.
45. **(New)** The method of claim 44, wherein the lymphoid cell is a Th2 cell.
46. **(New)** The method of claim 44, wherein the lymphoid cell is a Th1 cell.
47. **(New)** The method of claim 44, wherein the lymphoid cell is a B cell.
48. **(New)** The method of claim 43, wherein the indicator cell is a non-lymphoid mammalian cell.
49. **(New)** The method of claim 43, wherein the indicator cell is a yeast cell.
50. **(New)** A method for identifying a protein from Th2 cells that interacts with a maf family protein, comprising:
- a) providing a population of indicator cells, which population of indicator cells comprises:
    - i) a maf family protein lacking a transcriptional activation domain;
    - ii) a cDNA library prepared from Th2 cells; and
    - iii) a reporter gene operably linked to a transcriptional regulatory sequence to which the maf family protein binds;wherein expression of the reporter gene is sensitive to interactions between the maf family protein, a protein encoded by the cDNA library and the transcriptional regulatory sequence; and
  - b) selecting indicator cells within the population in which reporter gene expression is stimulated to thereby identify a protein from Th2 cells that interacts with a maf family protein.
51. **(New)** The method of claim 50, wherein the maf family protein is a c-Maf protein.
52. **(New)** The method of claim 50, wherein the maf family protein is selected from the group consisting of v-maf, mafB, Nrl, mafK, mafF, mafG and p18.

0924675-034260

53. (New) The method of claim 50, wherein the transcriptional regulatory sequence is from a Th2-associated cytokine gene.

54. (New) The method of claim 53, wherein the Th2-associated cytokine gene is an interleukin-4 gene.

55. (New) The method of claim 50, wherein the indicator cell is a yeast cell.--

### REMARKS

Claims 1-34 were present in the application as filed. Claims 1-34 have been canceled herein. Claims 35-55 have been added. Accordingly, claims 35-55, drawn to methods of identifying compounds that modulate the activity of a maf family protein or modulate an immune response, are currently pending in the application.


No new matter has been added. Support for the new claims can be found in originally filed claims 32-34 and throughout the specification, for example, page 36, line 4 through page 40, line 4. Support for new claims 40, 51, and 52 can be found at, for example, page 7, line 32 through page 8, line 1. Support for claims 44-48 can be found at, for example, page 37, lines 18-21. Support for new claim 50 can be found at, for example, page 39, lines 4-14.

Cancellation of and/or amendment to the claims should in no way be construed as an acquiescence to any of the Examiner's rejections. The cancellation of and/or amendments to the claims are being made solely to expedite prosecution of the above-identified application. Applicants reserve the option to further prosecute the same or similar claims in the instant or in another patent application.

SUMMARY

If a telephone conversation with Applicants' Agent would expedite the prosecution of the above-identified application, the examiner is urged to call Applicants' Agent at (617) 227-7400.

Respectfully submitted,



Megan E. Williams, Ph.D.

Registration No. 43,270

Agent for Applicants

LAHIVE & COCKFIELD, LLP  
28 State Street  
Boston, MA 02109  
Tel. (617) 227-7400

Dated: February 12, 1999

652720-95284260

Applicant or Patentee: Laurie H. Glimcher, I-Cheng Ho Attorney's  
Serial or Patent No.: 08/636,602 Docket No.: HUI-021  
Filed or Issued: April 23, 1996  
Title: METHODS AND COMPOSITIONS FOR REGULATING T CELL SUBSETS BY MODULATING  
TRANSCRIPTION FACTOR ACTIVITY

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS  
(37 CFR 1.9(f) and 1.27(c)) - SMALL BUSINESS CONCERN

I hereby declare that I am

- ☐ the owner of the small business concern identified below:  
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF SMALL BUSINESS CONCERN Boston Life Sciences, Inc.  
ADDRESS OF SMALL BUSINESS CONCERN Reservoir Place  
1601 Trapelo Road  
Waltham, MA 02154

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 CFR 121.12, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees to the United States Patent and Trademark Office, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention described in:

- ☐ the specification filed herewith with title as listed above.  
☒ the application identified above.  
☐ the patent identified above.

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights in the invention is listed below\* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

\*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME President and Fellows of Harvard College  
ADDRESS 124 Mt. Auburn Street, Cambridge, MA 02138  
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☒ NONPROFIT ORGANIZATION

NAME \_\_\_\_\_  
ADDRESS \_\_\_\_\_  
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING Marc Lanser

TITLE OF PERSON OTHER THAN OWNER Executive Vice President

ADDRESS OF PERSON SIGNING Boston Life Science, Inc., Reservoir Place, 1601 Trapelo Road, Waltham, MA 02154

SIGNATURE [Signature] DATE 6/17/96



Applicant or Patentee: Laurie H. Glimcher and I-Cheng Ho

Attorney's

Serial or Patent No.: 08/636,602

Docket No.: HUI-021

Filed or Issued: April 23, 1996

Title: METHODS AND COMPOSITIONS FOR REGULATING T CELL SUBSETS BY  
MODULATING TRANSCRIPTION FACTOR ACTIVITY

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS**  
**(37 CFR 1.9(f) and 1.27(d)) - NONPROFIT ORGANIZATION**

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF NONPROFIT ORGANIZATION President and Fellows of Harvard College

ADDRESS OF NONPROFIT ORGANIZATION 124 Mt. Auburn Street, Cambridge, MA 02138

TYPE OF NONPROFIT ORGANIZATION

- ☒ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION  
☐ TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 USC 501(a) and 501(c)(3))  
☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA  
(NAME OF STATE \_\_\_\_\_)  
(CITATION OF STATUTE \_\_\_\_\_)  
☐ WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 USC 501(a) and 501(c)(3)) IF LOCATED IN THE UNITED STATES OF AMERICA  
☐ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA  
(NAME OF STATE \_\_\_\_\_)  
(CITATION OF STATUTE \_\_\_\_\_)

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees to the United States Patent and Trademark Office regarding the invention entitled Methods and Compositions For Regulating T Cell Subsets By Modulating Transcription Factor Activity

by inventor(s) Laurie H. Glimcher and I-Cheng Ho

described in

- ☐ the specification filed herewith.  
☒ application serial no. 08/636,602, filed April 23, 1996  
☐ patent no. \_\_\_\_\_, issued \_\_\_\_\_

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization regarding the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights in the invention is listed below\* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

\*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME Boston Life Sciences, Inc.

ADDRESS Reservoir Place, 1601 Trapelo Road, Waltham, MA 02154

☐ INDIVIDUAL ☒ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING \_\_\_\_\_  
TITLE IN ORGANIZATION OF PERSON SIGNING \_\_\_\_\_  
ADDRESS OF PERSON SIGNING \_\_\_\_\_

Joyce Brinton, Director  
Office for Technology and Trademark Licensing  
Harvard University

SIGNATURE

DATE

**METHODS AND COMPOSITIONS FOR REGULATING T CELL SUBSETS BY  
MODULATING TRANSCRIPTION FACTOR ACTIVITY**

5

**Government Funding**

Work described herein was supported, at least in part, under grant AI37833 awarded by the National Institutes of Health. The U.S. government therefore may have certain rights in this invention.

10

**Background of the Invention**

CD4+ T helper cells are not a homogeneous population but can be divided on the basis of cytokine secretion into at least two subsets termed T helper type 1 (Th1) and T helper type 2 (Th2) (see *e.g.*, Mosmann, T.R. *et al.* (1986) *J. Immunol.* 136:2348-2357; Paul, W.E. and Seder, R.A. (1994) *Cell* 76:241-251; Seder, R.A. and Paul, W.E. (1994) *Ann. Rev. Immunol.* 12:635-673). Th1 cells secrete interleukin-2 (IL-2) and interferon- $\gamma$  (IFN- $\gamma$ ) while Th2 cells produce interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-10 (IL-10) and interleukin-13 (IL-13). Both subsets produce cytokines such as tumor necrosis factor (TNF) and granulocyte/macrophage-colony stimulating factor (GM-CSF). In addition to their different pattern of cytokine expression, Th1 and Th2 cells are thought to have differing functional activities. For example, Th1 cells are involved in inducing delayed type hypersensitivity responses, whereas Th2 cells are involved in providing efficient "help" to B lymphocytes and stimulating production of IgG1 and IgE antibodies.

There is now abundant evidence that the ratio of Th1 to Th2 cells is highly relevant to the outcome of a wide array of immunologically-mediated clinical diseases including autoimmune, allergic and infectious diseases. For example, in experimental leishmania infections in mice, animals that are resistant to infection mount predominantly a Th1 response, whereas animals that are susceptible to progressive infection mount predominantly a Th2 response (Heinzel, F.P., *et al.* (1989) *J. Exp. Med.* 169:59-72; Locksley, R. M. and Scott, P. (1992) *Immunoparasitology Today* 1:A58-A61). In murine schistosomiasis, a Th1 to Th2 switch is observed coincident with the release of eggs into the tissues by female parasites and is associated with a worsening of the disease condition (Pearce, E.J., *et al.* (1991) *J. Exp. Med.* 173:159-166; Grzych, J-M., *et al.* (1991) *J. Immunol.* 141:1322-1327; Kullberg, M.C., *et al.* (1992) *J. Immunol.* 148:3264-3270). Many human diseases, including chronic infections (such as with human immunodeficiency virus (HIV) and tuberculosis) and certain metastatic carcinomas, also are characterized by a Th1 to Th2 switch (see *e.g.*, Shearer, G.M. and Clerici, M. (1992) *Prog. Chem. Immunol.* 54:21-43; Clerici, M and Shearer, G.M. (1993) *Immunology Today* 14:107-111; Yamamura, M., *et al.* (1993) *J. Clin. Invest.* 91:1005-1010; Pisa, P., *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:7708-7712; Fauci,

09248756-DE-1299

A.S. (1988) *Science* 239:617-623). Furthermore, certain autoimmune diseases have been shown to be associated with a predominant Th1 response. For example, patients with rheumatoid arthritis have predominantly Th1 cells in synovial tissue (Simon, A.K., *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:8562-8566) and experimental autoimmune encephalomyelitis (EAE) can be induced by autoreactive Th1 cells (Kuchroo, V.K., *et al.* (1993) *J. Immunol.* 151:4371-4381).

The ability to alter or manipulate ratios of Th1 and Th2 subsets requires an understanding of the mechanisms by which the differentiation of CD4 T helper precursor cells (Thp), which secrete only IL-2, choose to become Th1 or Th2 effector cells. It is clear that the cytokines themselves are potent Th cell inducers and form an autoregulatory loop (see *e.g.*, Paul, W.E. and Seder, R.A. (1994) *Cell* 76:241-251; Seder, R.A. and Paul, W.E. (1994) *Ann. Rev. Immunol.* 12:635-673). Thus, IL-4 promotes the differentiation of Th2 cells while preventing the differentiation of precursors into Th1 cells, while IL-12 and IFN- $\gamma$  have the opposite effect. One possible means therefore to alter Th1:Th2 ratios is to increase or decrease the level of selected cytokines. Direct administration of cytokines or antibodies to cytokines has been shown to have an effect on certain diseases mediated by either Th1 or Th2 cells. For example, administration of recombinant IL-4 or antibodies to IL-12 ameliorate EAE, a Th1-driven autoimmune disease (see Racke; M.K. *et al.* (1994) *J. Exp. Med.* 180:1961-1966; and Leonard, J.P. *et al.* (1995) *J. Exp. Med.* 181:381-386), while anti-IL-4 antibodies cure the Th2-mediated parasitic disease, *Leishmania major* (Sadick, M.D. *et al.* (1990) *J. Exp. Med.* 171:115-127). However, as therapeutic options, systemic administration of cytokines or antibodies may have unwanted side effects and, accordingly, alternative approaches to manipulating Th1/Th2 subsets are still needed.

The molecular basis for the tissue-specific expression of IL-4 in Th2 cells, or any T cell cytokine, has remained elusive. One possibility is the presence of repressor proteins that selectively silence cytokines. Transcriptional silencing has been well documented for bacteria, yeast and mammalian genes. Examples include *E. coli* thermoregulation genes (Goransson, M. *et al.* (1990) *Nature* 344:682-685), yeast  $\alpha 2$  mating type genes (Keleher, C.A. *et al.* (1988) *Cell* 53:927-936) and mammalian MHC class I and TcR $\alpha$  genes (Weisman, J.D. and Singer, D.S. (1991) *Mol. Cell. Biol.* 11:4228-4234; Winoto, A. and Baltimore, D. (1989) *Cell* 59:649-655). Indeed, early experiments involving injection of IL-2 genomic DNA into *Xenopus* oocytes suggested the existence of a repressor protein for IL-2 in resting versus activated T cell extracts (Mouzaki, A. *et al.* (1991) *EMBO J.* 10:1399-1406). These studies suggested that the absence of IL-2 production in resting T cells was due to proteins that silenced the transcription of IL-2 by interacting with negative elements in the IL-2 promoter.

A second possibility is the existence of Th selective transactivators. A family of four related transcription factors called Nuclear Factor of Activated T cells (NF-AT), plays a key role in the regulation of cytokine gene expression (see *e.g.*, Emmel, E.A. *et al.* (1989) *Science*

246:1617-1620; Flanagan, W.M. *et al.* (1991) *Nature* 352:803-807; Jain, J. *et al.* (1993) *Nature* 365:352-355; McCaffrey, P.G. *et al.* (1993) *Science* 262:750-754; Rao, A. (1994) *Immunol. Today* 15:274-281; Northrop, J.P. *et al.* (1994) *Nature* 369:497). However, NF-AT family members can bind to and transactivate the promoters of multiple cytokine genes including IL-2 and IL-4 (Rooney, J. *et al.* (1995) *Immunity* 2:545-553; Szabo, S.J. *et al.* (1993) *Mol. Cell. Biol.* 13:4793-4805; Flanagan, W.M. *et al.* (1991) *Nature* 352:803-807; Northrop, J.P. *et al.* (1994) *Nature* 369:497). Thus, they are not likely to be responsible for directing Th1- or Th2-specific cytokine transcription. Most, if not all, NF-AT binding sites in cytokine promoter regulatory regions are accompanied by nearby sites that bind auxiliary transcription factors, usually members of the AP-1 family. It has been shown that NF-AT and AP-1 proteins bind coordinately and cooperatively and are required for full activity of the IL-2 and IL-4 promoters. Different AP-1 proteins, specifically c-Jun, c-Fos, Fra-1, Fra-2, Jun B and Jun D, have been shown to bind to these sites (Rao, A. *et al.* (1994) *Immunol. Today* 15:274-281; Jain, J. *et al.* (1993) *Nature* 365:352-355; Boise, L.H. *et al.* (1993) *Mol. Cell. Biol.* 13:1911-1919; Rooney, J. *et al.* (1995) *Immunity* 2:545-553; Rooney, J. *et al.* (1995) *Mol. Cell. Biol.* 15:6299-6310). However, none of these AP-1 proteins is expressed in a Th1- or Th2-specific manner and there is no evidence for the differential recruitment of AP-1 family members to the IL-2 or IL-4 composite sites (Rooney, J. *et al.* (1995) *Mol. Cell. Biol.* 15:6299-6310). Thus, neither NF-AT proteins nor the AP-1 family members c-Jun, c-Fos, Fra-1, Fra-2, Jun B and Jun D can account for the tissue-specific transcription of IL-4 in Th2 cells.

### **Summary of the Invention**

This invention pertains to methods for regulating Th1 or Th2 subsets by modulating the activity of a transcription factor that regulates expression of a Th2-specific cytokine gene. As described further herein, it has now been discovered that the tissue-specific expression of IL-4 in Th2 cells is not due to a repressor protein but rather to a Th2-specific transactivator protein. The proto-oncogene *c-maf* has now been demonstrated to be responsible for the tissue-specific expression of the Th2-associated cytokine interleukin-4. Moreover, ectopic expression of *c-maf* in cells other than Th2 cells (e.g, Th1 cells, B cells and non-lymphoid cells) leads to activation of the IL-4 promoter and, under appropriate conditions, production of endogenous IL-4.

Accordingly, in one aspect the invention provides a method for modulating production of a T helper type 2 (Th2)-associated cytokine by a cell. The method involves contacting the cell with an agent that modulates the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene such that production of the Th2-associated cytokine by a cell is modulated. In particular, the agents of the invention act intracellularly to modulate the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene. Preferably, the transcription factor is a member of the *maf*

family. Most preferably, the transcription factor is c-Maf. The Th2-associated cytokine modulated in the method is preferably interleukin-4. In one embodiment, production of the Th2-associated cytokine (*e.g.*, IL-4) is stimulated, for example in a cell that does not normally express the cytokine (such as a Th1 cell or a B cell). A variety of agents can be used to stimulate cytokine production, including a nucleic acid molecule encoding a maf family protein that is introduced into and expressed in the cell and chemical agents that enhance the expression or activity of an endogenous maf family protein in the cell. In another embodiment, production of a Th2-associated cytokine by a cell (*e.g.*, a Th2 cell) is inhibited. A variety of agents can be used to inhibit cytokine production, including antisense nucleic acid molecules that are complementary to a maf family gene, intracellular antibodies that bind maf family proteins (*e.g.*, in the cell nucleus), inhibitory forms of maf family proteins (*e.g.*, dominant negative forms) and chemical agents that inhibit the expression or activity of an endogenous maf family protein in the cell. Cytokine production by the cell can be modulated *in vitro* or *in vivo*. In one embodiment, a cell is contacted with a modulating agent *in vitro* and then is administered to a subject to thereby regulate the development of Th1 and/or Th2 subsets in the subject.

In another aspect, the invention provides methods for regulating the development of Th1 or Th2 subsets in a subject. In addition to the embodiment discussed above wherein *ex vivo* modified cells are administered to the subject, in another embodiment, these methods involve direct administration to the subject of an agent that modulates the activity of a transcription factor (*e.g.*, a maf family member) that regulates expression of a Th2-associated cytokine gene (*e.g.*, IL-4) such that development of Th1 or Th2 cells in the subject is modulated.

The methods of the invention can further involve the use of additional agents that modulate the activity of additional transcription factors that contribute to regulating the expression of Th1- or Th2-associated cytokines. Preferred additional agents are those which modulate the activity of a Nuclear Factor of Activated T cells (NF-AT) protein. Thus, in one embodiment, a stimulatory method of the invention can involve the use of a first agent that stimulates the expression and/or activity of a maf protein and a second agent that stimulates the expression and/or activity of an NF-AT protein. Similarly, an inhibitory method of the invention can involve the use of a first agent that inhibits the expression and/or activity of a maf protein and a second agent that inhibits the expression and/or activity of an NF-AT protein. Alternatively or additionally, the modulatory methods of the invention can involve the use of additional agents that modulate the activity of an AP-1 family protein.

The modulatory methods of the invention can be used to manipulate Th1:Th2 ratios in a variety of clinical situations. For example, inhibition of Th2 formation may be useful in allergic diseases, malignancies and infectious diseases whereas enhancement of Th2 formation may be useful in autoimmune diseases and organ transplantation.

Another aspect of the invention pertains to compositions that are useful for modulating the production of a Th2-associated cytokine by a cell and/or for modulating the development of Th1 or Th2 subsets in a subject. In one embodiment, these compositions include recombinant expression vectors that encode a maf family protein, wherein the maf-  
5 encoding sequences are operatively linked to regulatory sequences that direct expression of the maf family protein in a specific cell type, such as lymphoid cells (*e.g.*, T cells or B cells) or hematopoietic stem cells. In another embodiment, these compositions include host cells, such as host lymphoid cells (*e.g.*, host T cells or host B cells) or host hematopoietic stem cells, into which a recombinant expression vector encoding a maf family protein has been  
10 introduced.

Yet another aspect of the invention pertains to screening assays for identifying compounds that modulate the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene. In one type of screening assay, an indicator cell which contains both 1) a recombinant expression vector encoding a transcription factor that  
15 regulates expression of a Th2-associated cytokine gene and 2) a vector comprising regulatory sequences of the Th2-associated cytokine gene operatively linked a reporter gene is used to identify compounds that modulate the expression and/or activity of the transcription factor. In another embodiment, a screening assay of the invention identifies proteins from Th2 cells that form a protein-protein interaction with a transcription factor (*e.g.*, c-Maf) that regulates  
20 expression of a Th2-associated cytokine gene. In yet another embodiment, a screening assay of the invention identifies compounds that modulate the interaction of c-Maf with a maf response element (MARE) in the promoter of a Th2-associated cytokine gene (*e.g.*, a MARE in the IL-4 promoter).

## 25 **Brief Description of the Drawings**

*Figure 1A* is a schematic of the cell fusion approach used to demonstrate that cytokine expression is not due to a repressor

*Figure 1B* is a reverse transcriptase-polymerase chain reaction (RT-PCR) analysis of IL-2 and IL-4 cytokine, and control  $\beta$ -actin, mRNA expressed by an unfused Th1 clone  
30 (D1.1), an unfused Th2 clone (D10), Th1 and Th2 homokaryons and Th1-Th2 heterokaryons.

*Figure 2A* is a Northern blot analysis depicting expression of an isolated cDNA clone in Th1 cells, Th2 cells or B lymphoma cells. A control probe specific for GAPDH was used to show equal loading of RNA.

*Figure 2B* is a Northern blot analysis depicting upregulated expression of the isolated  
35 cDNA clone during *in vitro* differentiation of normal naive spleen cells into Th2 cells. Total RNA was isolated from cells harvested at the indicated time points. Culture supernatant at the appropriate dilution was measured for cytokine (IL-10) production by ELISA to determine differentiation into the Th1 or Th2 lineage.

*Figure 3A* is a bar graph depicting transactivation of the IL-4 promoter by c-Maf in a Th1 clone (AE7). AE7 cells were cotransfected with a wild-type IL-4 CAT reporter construct and either a control plasmid (pMEX-NeoI), a c-Maf expression plasmid (pMEX-Maf) or a c-Fos expression plasmid (pMEX-c-Fos). Half of each sample was stimulated 24 hours after transfection with antibodies to CD3. All samples were harvested 48 hours after transfection and relative CAT activities were determined.

*Figure 3B* is a photograph of a thin layer chromatography plate depicting the relative CAT activity in M12 B lymphoma cells cotransfected with a wild-type IL-4 CAT reporter construct and either two control plasmids (pMEX-NeoI and pREP4), a c-Maf expression plasmid and a control plasmid (pMEX-Maf and pREP4), a c-Fos expression plasmid and a control plasmid (pMEX-c-Fos and pREP4), a c-Jun expression plasmid and a control plasmid (pMEX-c-Jun and pREP4), a control plasmid and an NF-ATp expression plasmid (pMEX-NeoI and pREP-NFATp), a c-Maf expression plasmid and an NFATp expression plasmid (pMEX-Maf and pREP-NFATp) or a c-Fos expression plasmid and an NF-ATp expression plasmid (pMEX-c-Fos and pREP-NFATp). Half of each sample was stimulated 24 hours after transfection with PMA and ionomycin. All samples were harvested 48 hours after transfection and relative CAT activities were determined.

*Figure 4* is a bar graph depicting endogenous production of IL-4 in M12 cells by ectopic expression of c-Maf and NF-ATp. Cells stably transfected with the indicated control or expression plasmids were either unstimulated or stimulated with PMA and ionomycin for 24 hours. 200 µl of supernatant from each sample was subjected to ELISA for cytokine quantitation.

*Figure 5A* is a photograph of a DNase I footprint gel of the IL-4 promoter performed using nuclear extracts from Th2 (D10, CDC35) or Th1 (AE7, S53) clones harvested at the indicated time points after stimulation with anti-CD3 antibodies, which depicts a Th2-specific footprint immediately downstream of the putative MARE site in the IL-4 promoter. Two Th2-specific hypersensitive residues on the non-coding strand of the IL-4 promoter are indicated by \*. Five lanes of a DNase I digestion of the IL-4 promoter probe (without nuclear extract) and a Maxam-Gilbert A/G ladder were run next to the DNase I treated samples.

*Figure 5B* is a schematic representation of the proximal regulatory region of the murine IL-4 promoter. The top portion shows the primary sequence of the murine IL-4 promoter. The numbers indicated are relative to the start site of transcription at +1. Asterisks denote the Th2-specific hypersensitive residues seen on DNase I footprint. The bottom portion shows the sequence of the wild type (-59 to -28) oligonucleotide and the 4 bp mutants used in EMSA and the functional assays shown in Figures 6 and 7. Altered nucleotides are shown in lowercase bold and correspond to the numbering system shown in the top portion.

*Figure 6* is a photograph of an electrophoretic mobility shift assay (EMSA) demonstrating that c-Maf but not c-Jun binds to the proximal IL-4 promoter and forms a

complex with NF-ATp. EMSA was performed using the indicated proteins and labeled double-stranded oligonucleotides.

*Figure 7A* is a bar graph (top) and a photograph of a thin layer chromatography plate (bottom) depicting the relative CAT activity in M12 cells co-transfected with a c-Maf expression vector and either the wild-type IL-4 CAT reporter construct or one of the 4 bp mutants shown in Figure 5B, demonstrating that transactivation of the IL-4 promoter by c-Maf maps to the MARE and Th2-specific footprint. The average of three independent experiments and one representative experiment are shown in the top and bottom portions, respectively.

*Figure 7B* is a photograph of an EMSA, performed using recombinant c-Maf, the IL-4 promoter (-59 to -27) probe and the indicated unlabeled double-stranded oligonucleotides as competitors, demonstrating that binding of recombinant c-Maf to the IL-4 promoter maps to the MARE and Th2-specific footprint.

### 15 **Detailed Description of the Invention**

This invention pertains to methods and compositions for regulating T cell subsets by modulating transcription factor activity, as well as to screening assays for identifying compounds that can modulate such transcription factor activity. The invention is based, at least in part, on the discovery that Th2-specific expression of the interleukin-4 gene does not result from the action of a specific repressor protein (as shown in Example 1) but rather from the action of a specific transactivator protein. As described further herein, the transcription factor responsible for Th2-specific expression of the interleukin-4 gene has now been identified as the c-Maf proto-oncoprotein, which is selectively expressed in differentiating and mature Th2 cells and absent from Th1 cells (see Example 2). Ectopic expression of c-Maf in cells that do not normally express it (such as Th1 cells and B cells) leads to transactivation of the IL-4 promoter (see Example 3) and, under appropriate conditions, to production of endogenous IL-4 (see Example 4). Moreover, a protein present in nuclear extracts of Th2 cells, but not Th1 cells, footprints the IL-4 promoter in the region of a maf response element (MARE) (see Example 5) and recombinant c-Maf binds to the IL-4 promoter *in vitro* (see Example 6). The ability of c-Maf to transactivate IL-4 maps to the MARE and Th2-specific footprint in the IL-4 promoter (see Example 7).

The maf family of proteins is a sub-family of AP-1/CREB/ATF proteins. The *v-maf* oncogene was originally isolated from a spontaneous musculoaponeurotic fibrosarcoma of chicken and identified as the transforming gene of the avian retrovirus, AS42 (Nishizawa, M. *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:7711-7715). *V-maf* encodes a 42 kd basic region/leucine zipper (b-zip) protein with homology to the *c-fos* and *c-jun* oncogenes. Its cellular homologue, the *c-maf* proto-oncogene has only two structural changes in the coding region from v-maf (Kataoka, K. *et al.* (1993) *J. Virol.* 67:2133-2141). The maf family includes *c-maf*, *mafB*, a human retina-specific gene *Nrl* (Swaroop, A. *et al.* (1992) *Proc. Natl.*



Acad. Sci. USA 89:266-270), *mafK*, *mafF*, *mafG* and *p18*. The latter four, *mafK*, *mafF*, *mafG* and *p18*, each encode proteins that lack the amino terminal two thirds of c-Maf that contains the transactivating domain (Fujiwara, K.T. *et al.* (1993) *Oncogene* 8:2371-2380; Igarashi, K. *et al.* (1995) *J. Biol. Chem.* 270:7615-7624; Andrews, N.C. *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:11488-11492; Kataoka, K. *et al.* (1995) *Mol. Cell. Biol.* 15:2180-2190) and are referred to herein as "small" mafs. C-maf and maf family members form homodimers and heterodimers with each other and with Fos and Jun, consistent with the known ability of the AP-1 proteins to pair with each other (Kerppola, T.K. and Curran, T. (1994) *Oncogene* 9:675-684; Kataoka, K. *et al.* (1994) *Mol. Cell. Biol.* 14:700-712). The DNA target sequence to which c-Maf homodimers bind, termed the c-Maf response element (MARE), is a 13 or 14 bp element which contains a core TRE (T-MARE) or CRE (C-MARE) palindrome respectively. Prior to the present invention, little was known about the function of maf family members, although c-Maf has been shown to stimulate transcription from the Purkinje neuron-specific promoter L7 (Kurscher, C. and Morgan, J.I. (1994) *Mol. Cell. Biol.* 15:246-254) and Nrl has been shown to drive expression of the QR1 retina-specific gene (Swaroop, A. *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:266-270). The small mafs have been shown to function as repressors of  $\alpha$  and  $\beta$ -globin transcription when bound as homodimers but are essential as heterodimeric partners with the erythroid-specific factor p45NF-E2 to activate globin gene transcription (Kataoka, K. *et al.* (1995) *Mol. Cell. Biol.* 15:2180-2190; Igarashi, K. *et al.* (1994) *Nature* 367:568-572). MafK overexpression has been shown to induce erythroleukemia cell differentiation (Igarashi, K. *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92:7445-7449). However, prior to the present invention, there have been no reports implicating *c-maf* or maf family members in the regulation of genes expressed in lymphoid cells or in cytokine gene expression in any tissue.

Various aspects of the present invention are described in further detail in the following subsections.

#### I. Modulation of Th2-Associated Cytokine Production

As discussed above, the transcription factor responsible for the Th2-specific expression of the interleukin-4 gene has now been identified as the c-Maf proto-oncogene. Modulation of the expression and/or activity of c-Maf, therefore, provides a means to regulate the production of interleukin-4. Since IL-4 itself serves an autoregulatory function in the development of Th2 cells (see *e.g.*, Paul, W.E. and Seder, R.A. (1994) *Cell* 76:241-251; Seder, R.A. and Paul, W.E. (1994) *Ann. Rev. Immunol.* 12:635-673), and thus production of IL-4 can lead to the production of additional Th2-associated cytokines such as IL-5, IL-10 and IL-13 through further Th2 differentiation, modulation of c-Maf expression and/or activity provides a general approach for modulating production of Th2-associated cytokines.

Accordingly, the invention provides a method for modulating production of a Th2-associated cytokine by a cell. This method involves contacting the cell with an agent that

modulates the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene such that production of the Th2-associated cytokine by a cell is modulated. In particular, the modulatory agents of the invention are characterized by acting intracellularly to modulate the activity of a transcription factor. Preferably, the transcription factor is a member of the maf family and most preferably is c-Maf. As used herein, the term "Th2-associated cytokine" is intended to refer to a cytokine that is produced preferentially or exclusively by Th2 cells rather than by Th1 cells. Preferably the Th2-associated cytokine is interleukin-4. As used herein, the term "contacting" (*i.e.*, contacting a cell with an agent) is intended to include incubating the agent and the cell together *in vitro* (*e.g.*, adding the agent to cells in culture) and administering the agent to a subject such that the agent and cells of the subject are contacted *in vivo*. As used herein, the various forms of the term "modulation" are intended to include stimulation (*e.g.*, increasing or upregulating a particular response or activity) and inhibition (*e.g.*, decreasing or downregulating a particular response or activity). Accordingly, in one embodiment of the method of the invention, production of a Th2-associated cytokine by the cell is stimulated by contacting the cell with a stimulatory agent that stimulates c-Maf expression and/or activity. In another embodiment of the method of the invention, production of a Th2-associated cytokine by the cell is inhibited by contacting the cell with an inhibitory agent that inhibits c-Maf expression and/or activity.

As demonstrated in the Examples, although c-Maf is responsible for the tissue specificity of IL-4 gene expression, c-Maf acts synergistically with one or more additional transcription factors to activate IL-4 gene transcription. In particular, c-Maf acts synergistically with an NF-AT protein to stimulate IL-4 gene expression. Moreover, NF-AT proteins and other members of the AP-1/CREB/ATF family of transcription factors have been demonstrated to be involved in regulating expression of both Th1- and Th2-associated cytokine genes. Accordingly, the method of the invention for modulating Th2-associated cytokine production by a cell can further comprise contacting the cell with a second agent that modulates (*i.e.*, stimulates or inhibits) the expression or activity of a second transcription factor that contributes to regulating the expression of a Th1- or Th2-associated cytokine gene (discussed further below).

#### A. Stimulatory Agents

According to the method of the invention, to stimulate Th2-associated cytokine production by a cell, the cell is contacted with a stimulatory agent that stimulates expression and/or activity of a transcription factor (*e.g.*, c-Maf) that regulates expression of a Th2-associated cytokine gene. Th2-associated cytokine production can be stimulated in cell types that do not normally express such cytokines, such as Th1 cells, B cells or non-lymphoid cells.

Furthermore, Th2-associated cytokine production can be stimulated in helper precursor cells (Thp) to promote their differentiation along the Th2 pathway instead of the Th1 pathway.

A preferred stimulatory agent is a nucleic acid molecule encoding a maf family protein, wherein the nucleic acid molecule is introduced into the cell in a form suitable for expression of the maf family protein in the cell. For example, a c-Maf cDNA is cloned into a recombinant expression vector and the vector is transfected into the cell. As demonstrated in Example 3, ectopic expression of a *c-maf* recombinant expression vector in Th1 cells, B cells or non-lymphoid cells leads to activation of the IL-4 promoter. Additionally, under appropriate conditions (discussed in further detail below), transcription of the endogenous IL-4 gene is stimulated, leading to IL-4 production by cells that do not normally express this cytokine (see Example 4).

To express a maf family protein in a cell, typically a maf family cDNA is first introduced into a recombinant expression vector using standard molecular biology techniques. A maf family cDNA can be obtained, for example, by amplification using the polymerase chain reaction (PCR) or by screening an appropriate cDNA library. The nucleotide sequences of maf family cDNAs are known in the art and can be used for the design of PCR primers that allow for amplification of a cDNA by standard PCR methods or for the design of a hybridization probe that can be used to screen a cDNA library using standard hybridization methods. Preferably, the maf family cDNA is that of the *c-maf* proto-oncogene. The nucleotide and predicted amino acid sequences of a mammalian (mouse) *c-maf* cDNA are disclosed in Kurscher C. and Morgan, J.I. (1995) *Mol. Cell. Biol.* 15:246-254 and deposited in the GenBank database at accession number S74567. This mammalian *c-maf* is highly homologous to the avian *v-maf* sequence (disclosed in Nishizawa, M.K. *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:7711-7715 and GenBank accession numbers D28598 and D28596), indicating that *c-maf* is well conserved among species. *c-maf* cDNAs from other mammalian species, including humans, can be isolated using standard molecular biology techniques (*e.g.*, PCR or cDNA library screening) and primers or probes designed based upon the mouse or avian sequences. Human partial cDNA sequences homologous to the mouse c-maf cDNA are also deposited in the GenBank database at accession numbers H24189 and N75504. The sequences of other maf family members are also known in the art, for example MafB (Kataoka, K. *et al.* (1994) *Mol. Cell Biol.* 14:7581-91; GenBank accession number D28600), MafG (Kataoka *et al.* (1994) *Mol. Cell Biol.* 14:7581-91; GenBank accession numbers D28601 and D28602), MafF (GenBank accession number D16184) and MafK (Igarashi, K. *et al.* (1995) *J. Biol. Chem.* 270:7615-7624; GenBank accession numbers D16187 and D42124).

Following isolation or amplification of a maf family cDNA, the DNA fragment is introduced into an expression vector. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which

additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors  
5 (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" or simply "expression vectors". In general, expression vectors of utility in recombinant DNA  
10 techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

15 The recombinant expression vectors of the invention comprise a nucleic acid in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression and the level of expression desired, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant  
20 expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation  
25 signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell, those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences) or those which direct  
30 expression of the nucleotide sequence only under certain conditions (*e.g.*, inducible regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, *etc.* When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example,  
35 commonly used promoters are derived from polyoma virus, adenovirus, cytomegalovirus and Simian Virus 40. Non-limiting examples of mammalian expression vectors include pCDM8 (Seed, B., (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987), *EMBO J.* 6:187-195). A variety of mammalian expression vectors carrying different regulatory sequences are commercially available. For constitutive expression of the nucleic acid in a mammalian host

- cell, a preferred regulatory element is the cytomegalovirus promoter/enhancer. Moreover, inducible regulatory systems for use in mammalian cells are known in the art, for example systems in which gene expression is regulated by heavy metal ions (see *e.g.*, Mayo *et al.* (1982) *Cell* 29:99-108; Brinster *et al.* (1982) *Nature* 296:39-42; Searle *et al.* (1985) *Mol. Cell. Biol.* 5:1480-1489), heat shock (see *e.g.*, Nouer *et al.* (1991) in *Heat Shock Response*, e.d. Nouer, L., CRC, Boca Raton, FL, pp167-220), hormones (see *e.g.*, Lee *et al.* (1981) *Nature* 294:228-232; Hynes *et al.* (1981) *Proc. Natl. Acad. Sci. USA* 78:2038-2042; Klock *et al.* (1987) *Nature* 329:734-736; Israel & Kaufman (1989) *Nucl. Acids Res.* 17:2589-2604; and PCT Publication No. WO 93/23431), FK506-related molecules (see *e.g.*, PCT Publication No. WO 94/18317) or tetracyclines (Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. *et al.* (1995) *Science* 268:1766-1769; PCT Publication No. WO 94/29442; and PCT Publication No. WO 96/01313). Still further, many tissue-specific regulatory sequences are known in the art, including the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916) and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the  $\alpha$ -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).
- 25 Vector DNA can be introduced into mammalian cells via conventional transfection techniques. As used herein, the various forms of the term "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into mammalian host cells, including calcium phosphate co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for
- 30 transfecting host cells can be found in Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker may be introduced into a host cell on a separate vector from

that encoding a maf family protein or, more preferably, on the same vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

Another form of a stimulatory agent for stimulating expression of a Th2-associated cytokine in a cell is a chemical compound that stimulates the expression or activity of an endogenous maf family protein in the cell. Such compounds can be identified using screening assays that select for compounds that stimulate the expression or activity of a maf family protein. Examples of suitable screening assays are described in further detail in subsection V below.

In addition to use of an agent that stimulates the expression or activity of a maf family protein, the stimulatory methods of the invention can involve the use of a second agent that stimulates the expression or activity of a second transcription factor that contributes to regulating the expression of a Th1- or Th2-associated cytokine gene. In Example 4, it is shown that stimulation of the expression of endogenous IL-4 in M12 B lymphoma cells required the introduction into the cells of both a c-Maf expression vector and an NF-AT expression vector, thereby demonstrating that c-Maf and NF-AT act synergistically to activate IL-4 transcription, with c-maf responsible for the tissue-specificity of expression. While the skilled artisan will appreciate that certain cells may express sufficient amounts of endogenous NF-AT such that use of a second agent that stimulates the expression or activity of NF-AT is unnecessary, in certain situations and with certain cell types it may be necessary to stimulate both c-Maf and NF-AT to achieve IL-4 production.

Accordingly, in one embodiment, the stimulatory methods of the invention involve the use of a first agent that stimulates the expression or activity of c-Maf and a second agent that stimulates the expression or activity of an NF-AT protein. A preferred second agent for stimulating NF-AT activity in a cell is a recombinant expression encoding an NF-AT, wherein the recombinant expression vector is introduced into the cell and NF-AT is expressed in the cell. NF-AT-encoding expression vectors can be prepared and introduced into cells as described above for c-Maf expression vectors. The nucleotide sequences of NF-AT cDNAs are known in the art and can be used for the design of PCR primers that allow for amplification of a cDNA by standard PCR methods or for the design of a hybridization probe that can be used to screen a cDNA library using standard hybridization methods. Four NF-AT family members have been identified (see e.g., Emmel, E.A. *et al.* (1989) *Science* 246:1617-1620; Flanagan, W.M. *et al.* (1991) *Nature* 352:803-807; Jain, J. *et al.* (1993) *Nature* 365:352-355; McCaffrey, P.G. *et al.* (1993) *Science* 262:750-754; Rao, A. (1994) *Immunol. Today* 15:274-281; Northrop, J.P. *et al.* (1994) *Nature* 369:497). Preferably, the NF-AT cDNA is that of NF-ATp. The nucleotide and predicted amino acid sequences of a mammalian NF-ATp cDNA are disclosed in McCaffrey, P.G. *et al.* (1993) *Science* 262:750-754. The nucleotide and predicted amino acid sequences of a mammalian NF-ATc cDNA are disclosed in Northrop, J.P. *et al.* (1994) *Nature* 369:497 and deposited in the GenBank

database at accession number U08015. The nucleotide and predicted amino acid sequences of mammalian NF-AT3 and NF-AT4 cDNAs are disclosed in Hoey, T. et al. (1995) *Immunity* 2:461-472.

Alternative to use of an NF-AT cDNA to stimulate the activity of NF-AT in a cell, one or more chemical compounds that stimulate NF-AT activity in a cell can be used as a second agent in a stimulatory method of the invention. Compounds that stimulate NF-AT activity in cells are known in the art (for a review see Rao, A. (1994) *Immunol. Today* 15:274-281). For example, stimulation of certain cells with the phorbol ester phorbol myristate acetate (PMA) and a calcium ionophore (e.g., ionomycin) results in translocation of NF-ATs to the cell nucleus (see e.g., Flanagan, W.M. et al. (1991) *Nature* 352:803-807; Jain, J. et al. (1993) *Nature* 365:352-355). Additionally, stimulation of T cells through the T cell receptor (TcR), for example with an anti-CD3 antibody, results in activation of NF-AT in the T cells.

In addition to NF-AT proteins, AP-1 family members, including c-Jun, c-Fos, Fra-1, Fra-2, Jun B and Jun D, have been shown to be involved in regulating the expression of both Th1- and Th2-associated cytokine genes (e.g., IL-2 and IL-4) (see e.g., Rao, A. et al. (1994) *Immunol. Today* 15:274-281; Jain, J. et al. (1993) *Nature* 365:352-355; Boise, L.H. et al. (1993) *Mol. Cell. Biol.* 13:1911-1919; Rooney, J. et al. (1995) *Immunity* 2:545-553; Rooney, J. et al. (1995) *Mol. Cell. Biol.* 15:6299-6310). Although these factors are not responsible for the Th1/Th2 specificity of expression of the cytokine genes, and these factors do not appear to synergize with c-Maf in regulating IL-4 gene expression (see the Examples), AP-1 family members have been shown to increase IL-4 expression in Th2 cells (see e.g., Rooney, J. et al. (1995) *Immunity* 2:545-553). Accordingly, in certain circumstances it may be beneficial, in addition to stimulating c-Maf activity (and possibly NF-AT activity), also to stimulate the activity of an AP-1 protein. Accordingly, in one embodiment, the stimulatory methods of the invention involve the use of a first agent that stimulates the expression or activity of c-Maf and a second agent that stimulates the expression or activity of an AP-1 protein. In another embodiment, the invention involves the use of a first agent that stimulates the expression or activity of c-Maf, a second agent that stimulates the expression or activity of an NF-AT protein and a third agent that stimulates the expression or activity of an AP-1 protein.

A preferred agent for stimulating AP-1 activity in a cell is a recombinant expression encoding an AP-1 protein, wherein the recombinant expression vector is introduced into the cell and the AP-1 protein is expressed in the cell. AP-1-encoding expression vectors can be prepared and introduced into cells as described above for c-Maf expression vectors. The nucleotide sequences of AP-1 cDNAs are known in the art. For example, the nucleotide and predicted amino acid sequences of human *c-fos* are disclosed in van Straaten, F. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:3183-3187. The nucleotide and predicted amino acid sequences of human *c-jun* are disclosed in Bohmann, D. et al. (1987) *Science* 238:1386-1392. The nucleotide and predicted amino acid sequences of human *jun-B* and *jun-D* are disclosed

09248756 "021299

in Nomura, N. *et al.* (1990) *Nucl. Acids Res.* 18:3047-3048. The nucleotide and predicted amino acid sequences of human *fra-1* and *fra-2* are disclosed in Matsui, M. *et al.* (1990) *Oncogene* 5:249-255. These sequences can be used for the design of PCR primers that allow for amplification of a cDNA by standard PCR methods or for the design of a hybridization probe that can be used to screen a cDNA library using standard hybridization methods. Alternatively, one or more chemical compounds that stimulate AP-1 activity in a cell can be used as additional agents in a stimulatory method of the invention. Compounds that stimulate AP-1 activity in cells are known in the art, including PMA/calcium ionophore (*e.g.*, ionomycin) and anti-CD3 antibodies.

#### B. Inhibitory Agents

According to the method of the invention, to inhibit Th2-associated cytokine production by a cell, the cell is contacted with an inhibitory agent that inhibits expression and/or activity of a transcription factor (*e.g.*, c-Maf) that regulates expression of a Th2-associated cytokine gene. Th2-associated cytokine production can be inhibited in, for example, Th2 cells or in helper precursor cells (Thp) to promote their differentiation along the Th1 pathway instead of the Th2 pathway. Inhibitory agents of the invention can be, for example, intracellular binding molecules that act to inhibit the expression or activity of the transcription factor. As used herein, the term "intracellular binding molecule" is intended to include molecules that act intracellularly to inhibit the expression or activity of a protein by binding to the protein or to a nucleic acid (*e.g.*, an mRNA molecule) that encodes the protein. Examples of intracellular binding molecules, described in further detail below, include antisense nucleic acids, intracellular antibodies and dominant negative inhibitors.

In one embodiment, an inhibitory agent of the invention is an antisense nucleic acid molecule that is complementary to a gene encoding a maf family protein, or to a portion of said gene, or a recombinant expression vector encoding said antisense nucleic acid molecule. The use of antisense nucleic acids to downregulate the expression of a particular protein in a cell is well known in the art (see *e.g.*, Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986; Askari, F.K. and McDonnell, W.M. (1996) *N. Eng. J. Med.* 334:316-318; Bennett, M.R. and Schwartz, S.M. (1995) *Circulation* 92:1981-1993; Mercola, D. and Cohen, J.S. (1995) *Cancer Gene Ther.* 2:47-59; Rossi, J.J. (1995) *Br. Med. Bull.* 51:217-225; Wagner, R.W. (1994) *Nature* 372:333-335). An antisense nucleic acid molecule comprises a nucleotide sequence that is complementary to the coding strand of another nucleic acid molecule (*e.g.*, an mRNA sequence) and accordingly is capable of hydrogen bonding to the coding strand of the other nucleic acid molecule. Antisense sequences complementary to a sequence of an mRNA can be complementary to a sequence found in the coding region of the mRNA, the 5' or 3' untranslated region of the mRNA or a region bridging the coding region and an untranslated



region (e.g., at the junction of the 5' untranslated region and the coding region). Furthermore, an antisense nucleic acid can be complementary in sequence to a regulatory region of the gene encoding the mRNA, for instance a transcription initiation sequence or regulatory element. Preferably, an antisense nucleic acid is designed so as to be complementary to a region preceding or spanning the initiation codon on the coding strand or in the 3' untranslated region of an mRNA. An antisense nucleic acid for inhibiting the expression of a Maf family protein in a cell can be designed based upon the nucleotide sequence encoding the Maf family protein, constructed according to the rules of Watson and Crick base pairing.

An antisense nucleic acid can exist in a variety of different forms. For example, the antisense nucleic acid can be an oligonucleotide that is complementary to only a portion of a maf family gene. An antisense oligonucleotides can be constructed using chemical synthesis procedures known in the art. An antisense oligonucleotide can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g. phosphorothioate derivatives and acridine substituted nucleotides can be used. To inhibit Maf protein expression in cells in culture, one or more antisense oligonucleotides can be added to cells in culture media, typically at 200 µg oligonucleotide/ml.

Alternatively, an antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., nucleic acid transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest). Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the expression of the antisense RNA molecule in a cell of interest, for instance promoters and/or enhancers or other regulatory sequences can be chosen which direct constitutive, tissue specific or inducible expression of antisense RNA. The antisense expression vector is prepared as described above for recombinant expression vectors, except that the cDNA (or portion thereof) is cloned into the vector in the antisense orientation. The antisense expression vector can be in the form of, for example, a recombinant plasmid, phagemid or attenuated virus. The antisense expression vector is introduced into cells using a standard transfection technique, as described above for recombinant expression vectors.

In another embodiment, an antisense nucleic acid for use as an inhibitory agent is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region (for reviews on ribozymes see e.g., Ohkawa, J. *et al.* (1995) *J. Biochem.* 118:251-258; Sigurdsson, S.T. and Eckstein, F. (1995) *Trends Biotechnol.* 13:286-289; Rossi, J.J. (1995) *Trends Biotechnol.* 13:301-306; Kiehnopf, M. *et al.* (1995) *J. Mol. Med.* 73:65-71). A ribozyme having specificity for a maf family mRNA can be designed based upon the nucleotide sequence of the maf family, preferably *c-maf*. For example, a derivative of a

*Tetrahymena* L-19 IVS RNA can be constructed in which the base sequence of the active site is complementary to the base sequence to be cleaved in a *c-maf* mRNA. See for example U.S. Patent Nos. 4,987,071 and 5,116,742, both by Cech *et al.* Alternatively, *c-maf* mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See for example Bartel, D. and Szostak, J.W. (1993) *Science* 261: 1411-1418.

Another type of inhibitory agent that can be used to inhibit the expression and/or activity of a Maf protein in a cell is an intracellular antibody specific for the Maf protein. The use of intracellular antibodies to inhibit protein function in a cell is known in the art (see *e.g.*, Carlson, J. R. (1988) *Mol. Cell. Biol.* 8:2638-2646; Biocca, S. *et al.* (1990) *EMBO J.* 9:101-108; Werge, T.M. *et al.* (1990) *FEBS Letters* 274:193-198; Carlson, J.R. (1993) *Proc. Natl. Acad. Sci. USA* 90:7427-7428; Marasco, W.A. *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:7889-7893; Biocca, S. *et al.* (1994) *Bio/Technology* 12:396-399; Chen, S-Y. *et al.* (1994) *Human Gene Therapy* 5:595-601; Duan, L *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:5075-5079; Chen, S-Y. *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:5932-5936; Beerli, R.R. *et al.* (1994) *J. Biol. Chem.* 269:23931-23936; Beerli, R.R. *et al.* (1994) *Biochem. Biophys. Res. Commun.* 204:666-672; Mhashilkar, A.M. *et al.* (1995) *EMBO J.* 14:1542-1551; Richardson, J.H. *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92:3137-3141; PCT Publication No. WO 94/02610 by Marasco *et al.*; and PCT Publication No. WO 95/03832 by Duan *et al.*).

To inhibit protein activity using an intracellular antibody, a recombinant expression vector is prepared which encodes the antibody chains in a form such that, upon introduction of the vector into a cell, the antibody chains are expressed as a functional antibody in an intracellular compartment of the cell. For inhibition of transcription factor activity according to the inhibitory methods of the invention, preferably an intracellular antibody that specifically binds the transcription factor is expressed within the nucleus of the cell. Nuclear expression of an intracellular antibody can be accomplished by removing from the antibody light and heavy chain genes those nucleotide sequences that encode the N-terminal hydrophobic leader sequences and adding nucleotide sequences encoding a nuclear localization signal at either the N- or C-terminus of the light and heavy chain genes (see *e.g.*, Biocca, S. *et al.* (1990) *EMBO J.* 9:101-108; Mhashilkar, A. M. *et al.* (1995) *EMBO J.* 14:1542-1551). A preferred nuclear localization signal to be used for nuclear targeting of the intracellular antibody chains is the nuclear localization signal of SV40 Large T antigen (see Biocca, S. *et al.* (1990) *EMBO J.* 9:101-108; Mhashilkar, A. M. *et al.* (1995) *EMBO J.* 14:1542-1551).

To prepare an intracellular antibody expression vector, antibody light and heavy chain cDNAs encoding antibody chains specific for the target protein of interest, *e.g.*, a Maf family protein, are isolated, typically from a hybridoma that secretes a monoclonal antibody specific for the maf protein. Preparation of antisera against Maf family proteins has been described in the art (see *e.g.*, Fujiwara, K.T. *et al.* (1993) *Oncogene* 8:2371-2380; Kataoka, K. *et al.* (1993) *J. Virol.* 67:2133-2141; Kerppola, T.K. and Curran, T. (1994) *Oncogene* 9:675-684; Igarashi, K *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92:7445-7449). Anti-Maf protein

antibodies can be prepared by immunizing a suitable subject, (e.g., rabbit, goat, mouse or other mammal) with a Maf protein immunogen. An appropriate immunogenic preparation can contain, for examples, recombinantly expressed Maf protein or a chemically synthesized Maf peptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975, *Nature* 256:495-497) (see also, Brown *et al.* (1981) *J. Immunol* 127:539-46; Brown *et al.* (1980) *J Biol Chem* 255:4980-83; Yeh *et al.* (1976) *PNAS* 76:2927-31; and Yeh *et al.* (1982) *Int. J. Cancer* 29:269-75). The technology for producing monoclonal antibody hybridomas is well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*, 54:387-402; M. L. Gefter *et al.* (1977) *Somatic Cell Genet.*, 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with a maf protein immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds specifically to the Maf protein. Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating an anti-Maf protein monoclonal antibody (see, e.g., G. Galfre *et al.* (1977) *Nature* 266:550-52; Gefter *et al.* *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinary skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines may be used as a fusion partner according to standard techniques, e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from the American Type Culture Collection (ATCC), Rockville, Md. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody that specifically binds the maf protein are identified by screening the hybridoma culture supernatants for such antibodies, e.g., using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal anti-maf antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with a maf protein or peptide to thereby isolate immunoglobulin library members that bind specifically to a Maf protein. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP<sup>TM</sup> Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* International Publication No. WO 92/18619; Dower *et al.* International Publication No. WO 91/17271; Winter *et al.* International Publication WO 92/20791; Markland *et al.* International Publication No. WO 92/15679; Breitling *et al.* International Publication WO 93/01288; McCafferty *et al.* International Publication No. WO 92/01047; Garrard *et al.* International Publication No. WO 92/09690; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clarkson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; Barbas *et al.* (1991) *PNAS* 88:7978-7982; and McCafferty *et al.* *Nature* (1990) 348:552-554.

Once a monoclonal antibody specific for the Maf protein has been identified (e.g., either a hybridoma-derived monoclonal antibody or a recombinant antibody from a combinatorial library), DNAs encoding the light and heavy chains of the monoclonal antibody are isolated by standard molecular biology techniques. For hybridoma derived antibodies, light and heavy chain cDNAs can be obtained, for example, by PCR amplification or cDNA library screening. For recombinant antibodies, such as from a phage display library, cDNA encoding the light and heavy chains can be recovered from the display package (e.g., phage) isolated during the library screening process. Nucleotide sequences of antibody light and heavy chain genes from which PCR primers or cDNA library probes can be prepared are known in the art. For example, many such sequences are disclosed in Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242 and in the "Vbase" human germline sequence database.

Once obtained, the antibody light and heavy chain sequences are cloned into a recombinant expression vector using standard methods. As discussed above, the sequences encoding the hydrophobic leaders of the light and heavy chains are removed and sequences encoding a nuclear localization signal (e.g., from SV40 Large T antigen) are linked in-frame to sequences encoding either the amino- or carboxy terminus of both the light and heavy chains. The expression vector can encode an intracellular antibody in one of several different

forms. For example, in one embodiment, the vector encodes full-length antibody light and heavy chains such that a full-length antibody is expressed intracellularly. In another embodiment, the vector encodes a full-length light chain but only the VH/CH1 region of the heavy chain such that a Fab fragment is expressed intracellularly. In the most preferred  
5 embodiment, the vector encodes a single chain antibody (scFv) wherein the variable regions of the light and heavy chains are linked by a flexible peptide linker (*e.g.*, (Gly<sub>4</sub>Ser)<sub>3</sub>) and expressed as a single chain molecule. To inhibit transcription factor activity in a cell, the expression vector encoding the transcription factor-specific intracellular antibody is introduced into the cell by standard transfection methods, as discussed hereinbefore.

10 Yet another form of an inhibitory agent of the invention is an inhibitory form of a Maf protein, also referred to herein as a dominant negative inhibitor. The maf family of proteins are known to homodimerize and to heterodimerize with other AP-1 family members, such as Fos and Jun (see *e.g.*, Kerppola, T.K. and Curran, T. (1994) *Oncogene* 9:675-684; Kataoka, K. *et al.* (1994) *Mol. Cell. Biol.* 14:700-712). One means to inhibit the activity of  
15 transcription factors that form dimers is through the use of a dominant negative inhibitor that has the ability to dimerize with functional transcription factors but that lacks the ability to activate transcription (see *e.g.*, Petrak, D. *et al.* (1994) *J. Immunol.* 153:2046-2051). By dimerizing with functional transcription factors, such dominant negative inhibitors can inhibit their functional activity. This process may occur naturally as a means to regulate gene  
20 expression. For example, there are a number of "small" maf proteins, such as *mafK*, *mafF*, *mafG* and *p18*, which lack the amino terminal two thirds of c-Maf that contains the transactivating domain (Fujiwara, K.T. *et al.* (1993) *Oncogene* 8:2371-2380; Igarashi, K. *et al.* (1995) *J. Biol. Chem.* 270:7615-7624; Andrews, N.C. *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:11488-11492; Kataoka, K. *et al.* (1995) *Mol. Cell. Biol.* 15:2180-2190).  
25 Homodimers of the small maf proteins act as negative regulators of transcription (Igarashi, K. *et al.* (1994) *Nature* 367:568-572) and three of the small maf proteins (MafK, MafF and MafG) have been shown to competitively inhibit transactivation mediated by the v-Maf oncoprotein (Kataoka, K. *et al.* (1996) *Oncogene* 12:53-62). Additionally, MafB has been identified as an interaction partner of Ets-1 and shown to inhibit Ets-1-mediated  
30 transactivation of the transferrin receptor and to inhibit erythroid differentiation (Sieweke, M.H. *et al.* (1996) *Cell* 85:49-60).

Accordingly, an inhibitory agent of the invention can be a form of a Maf protein that has the ability to dimerize with c-Maf but that lacks the ability to activate transcription. This dominant negative form of a Maf protein may be, for example, a small Maf protein (*e.g.*,  
35 MafK, MafF, MafG) that naturally lacks a transactivation domain, MafB or a mutated form of c-Maf in which the transactivation domain has been removed. Such dominant negative Maf proteins can be expressed in cells using a recombinant expression vector encoding the Maf protein, which is introduced into the cell by standard transfection methods. To express a mutant form of c-Maf lacking a transactivation domain, nucleotide sequences encoding the

amino terminal transactivation domain of c-Maf are removed from the *c-maf* cDNA by standard recombinant DNA techniques. Preferably, at least amino acids 1-122 are removed. More preferably, at least amino acids 1-187, or amino acids 1-257, are removed. Nucleotide sequences encoding the basic-leucine zipper region are maintained. The truncated cDNA is inserted into a recombinant expression vector, which is then introduced into a cell to allow for expression of the truncated c-maf, lacking a transactivation domain, in the cell.

Yet another type of inhibitory agent that can be used to inhibit the expression and/or activity of a maf protein in a cell is chemical compound that inhibits the expression or activity of an endogenous maf family protein in the cell. Such compounds can be identified using screening assays that select for compounds that inhibit the expression or activity of a maf family protein. Examples of suitable screening assays are described in further detail in subsection V below.

As discussed above with regard to stimulatory agents, the inhibitory methods of the invention can involve the use of one or more additional inhibitory agents that inhibit the expression or activity of one or more additional transcription factors that contributes to regulating the expression of a Th1- or Th2-associated cytokine gene. For example, in one embodiment, the inhibitory method of the invention comprises contacting a cell with a first agent that inhibits the expression or activity a maf family protein and a second agent that inhibits the expression or activity of an NF-AT protein. In another embodiment, the inhibitory method of the invention comprises contacting a cell with a first agent that inhibits the expression or activity a maf family protein and a second agent that inhibits the expression or activity of an AP-1 protein. In yet another embodiment, the inhibitory method of the invention comprises contacting a cell with a first agent that inhibits the expression or activity a maf family protein, a second agent that inhibits the expression or activity of an NF-AT protein and a third agent that inhibits the expression or activity of an AP-1 protein. Examples of types of inhibitory agents for inhibiting NF-AT or AP-1 proteins include antisense nucleic acids, intracellular antibodies, dominant negative inhibitors and chemical compounds that inhibit the endogenous proteins, as described above. Regarding the latter, it is known in the art that the nuclear translocation of NF-ATp is inhibited by the immunosuppressive drugs cyclosporin A and FK506 (see e.g., Rao, A. (1994) *Immunol. Today* 15:274-281; Rao, A. (1995) *J. Leukoc. Biol.* 57:536-542). Accordingly, in one embodiment of the inhibitory method, an immunosuppressive drug such as cyclosporin A or FK506 (or other related drug that inhibits the calcineurin pathway) is used in combination with an agent that inhibits the expression or activity of c-Maf.

The method of the invention for modulating production of Th2-associated cytokines by a cell can be practiced either *in vitro* or *in vivo* (the latter is discussed further in the following subsection). For practicing the method *in vitro*, cells can be obtained from a subject by standard methods and incubated (*i.e.*, cultured) *in vitro* with a stimulatory or

inhibitory agent of the invention to stimulate or inhibit, respectively, the production of Th2-associated cytokines. For example, peripheral blood mononuclear cells (PBMCs) can be obtained from a subject and isolated by density gradient centrifugation, *e.g.*, with Ficoll/Hypaque. Specific cell populations can be depleted or enriched using standard methods. For example, monocytes/macrophages can be isolated by adherence on plastic. T cells or B cells can be enriched or depleted, for example, by positive and/or negative selection using antibodies to T cell or B cell surface markers, for example by incubating cells with a specific primary monoclonal antibody (mAb), followed by isolation of cells that bind the mAb using magnetic beads coated with a secondary antibody that binds the primary mAb. Peripheral blood or bone marrow derived hematopoietic stem cells can be isolated by similar techniques using stem cell-specific mAbs (*e.g.*, anti-CD34 mAbs). Specific cell populations can also be isolated by fluorescence activated cell sorting according to standard methods. Monoclonal antibodies to cell-specific surface markers known in the art and many are commercially available.

When a stimulatory agent is used *in vitro*, resulting in stimulation of the production of Th2-associated cytokines, in particular IL-4, the cytokine(s) can be recovered from the culture supernatant for further use. For example, the culture supernatant, or a purified fraction thereof, can be applied to T cells in culture to influence the development of Th1 or Th2 cells *in vitro*. Alternatively, the culture supernatant, or a purified fraction thereof, can be administered to a subject to influence the development of Th1 vs. Th2 responses in the subject.

Moreover, cells treated *in vitro* with either a stimulatory or inhibitory agent can be administered to a subject to influence the development of a Th1 vs. Th2 response in the subject. Accordingly, in another embodiment, the method of the invention for modulating the production of Th2-associated cytokines by a cell further comprises administering the cell to a subject to thereby modulate development of Th1 or Th2 cells in a subject. Preferred cell types for *ex vivo* modification and readministration include T cells, B cells and hematopoietic stem cells. For administration to a subject, it may be preferable to first remove residual agents in the culture from the cells before administering them to the subject. This can be done for example by a Ficoll/Hypaque gradient centrifugation of the cells. For further discussion of *ex vivo* genetic modification of cells followed by readministration to a subject, see also U.S. Patent No. 5,399,346 by W.F. Anderson *et al.*

## II. Methods for Modulating Development of Th1 or Th2 Cells in a Subject

Another aspect of the invention pertains to a method for modulating development of Th1 or Th2 cells in a subject. The term "subject" is intended to include living organisms in which an immune response can be elicited. Preferred subjects are mammals. Examples of subjects include humans, monkeys, dogs, cats, mice, rats, cows, horses, goats and sheep. As discussed above, one way to modulate Th1/Th2 ratios in a subject is to treat cells (*e.g.*, T

cells, B cells or hematopoietic stem cells) *ex vivo* with one or more modulatory agents of the invention, such that production of a Th2-associated cytokine by the cells is modulated, followed by administration of the cells to the subject. In another embodiment, Th1/Th2 ratios are modulated in a subject by administering to the subject an agent that modulates the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene such that development of Th1 or Th2 cells in the subject is modulated. Preferably, the transcription factor is a maf family protein and most preferably a c-Maf protein. Preferably, the Th2-associated cytokine is IL-4. Development of a Th2 response in the subject can be promoted by administration of one or more stimulatory agents of the invention, whereas development of a Th1 response in the subject can be promoted by administration of one or more inhibitory agents of the invention. As discussed above, in certain situations it may be desirable, in addition to modulating the activity of a maf family protein, to also modulate the activity of other transcription factors involved in regulating Th1- or Th2-associated cytokine genes. Preferably, the activity of an NF-AT protein(s) is also modulated. Additionally or alternatively, the activity of an AP-1 protein(s) is also modulated.

For stimulatory or inhibitory agents that comprise nucleic acids (including recombinant expression vectors encoding transcription factors, antisense RNA, intracellular antibodies or dominant negative inhibitors), the agents can be introduced into cells of the subject using methods known in the art for introducing nucleic acid (*e.g.*, DNA) into cells *in vivo*. Examples of such methods include:

*Direct Injection:* Naked DNA can be introduced into cells *in vivo* by directly injecting the DNA into the cells (see *e.g.*, Acsadi *et al.* (1991) *Nature* 332:815-818; Wolff *et al.* (1990) *Science* 247:1465-1468). For example, a delivery apparatus (*e.g.*, a "gene gun") for injecting DNA into cells *in vivo* can be used. Such an apparatus is commercially available (*e.g.*, from BioRad).

*Receptor-Mediated DNA Uptake:* Naked DNA can also be introduced into cells *in vivo* by complexing the DNA to a cation, such as polylysine, which is coupled to a ligand for a cell-surface receptor (see for example Wu, G. and Wu, C.H. (1988) *J. Biol. Chem.* 263:14621; Wilson *et al.* (1992) *J. Biol. Chem.* 267:963-967; and U.S. Patent No. 5,166,320). Binding of the DNA-ligand complex to the receptor facilitates uptake of the DNA by receptor-mediated endocytosis. A DNA-ligand complex linked to adenovirus capsids which naturally disrupt endosomes, thereby releasing material into the cytoplasm can be used to avoid degradation of the complex by intracellular lysosomes (see for example Curiel *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:8850; Cristiano *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:2122-2126).

*Retroviruses:* Defective retroviruses are well characterized for use in gene transfer for gene therapy purposes (for a review see Miller, A.D. (1990) *Blood* 76:271). A recombinant retrovirus can be constructed having a nucleotide sequences of interest incorporated into the retroviral genome. Additionally, portions of the retroviral genome can be removed to render



the retrovirus replication defective. The replication defective retrovirus is then packaged into virions which can be used to infect a target cell through the use of a helper virus by standard techniques. Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology,

- 5 Ausubel, F.M. *et al.* (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines include  $\psi$ Crip,  $\psi$ Cre,  $\psi$ 2 and  $\psi$ Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells,
- 10 endothelial cells, lymphocytes, myoblasts, hepatocytes, bone marrow cells, *in vitro* and/or *in vivo* (see for example Eglitis, *et al.* (1985) *Science* 230:1395-1398; Danos and Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:6460-6464; Wilson *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:6141-6145; Huber *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:8039-8043; Ferry *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:8377-8381; Chowdhury *et al.* (1991) *Science* 254:1802-1805; van
- 15 Beusechem *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:7640-7644; Kay *et al.* (1992) *Human Gene Therapy* 3:641-647; Dai *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Hwu *et al.* (1993) *J. Immunol.* 150:4104-4115; U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT
- 20 Application WO 89/05345; and PCT Application WO 92/07573). Retroviral vectors require target cell division in order for the retroviral genome (and foreign nucleic acid inserted into it) to be integrated into the host genome to stably introduce nucleic acid into the cell. Thus, it may be necessary to stimulate replication of the target cell.

- Adenoviruses:* The genome of an adenovirus can be manipulated such that it encodes
- 25 and expresses a gene product of interest but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. See for example Berkner *et al.* (1988) *BioTechniques* 6:616; Rosenfeld *et al.* (1991) *Science* 252:431-434; and Rosenfeld *et al.* (1992) *Cell* 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (*e.g.*, Ad2, Ad3, Ad7 *etc.*) are well known to those skilled in the art.
- 30 Recombinant adenoviruses are advantageous in that they do not require dividing cells to be effective gene delivery vehicles and can be used to infect a wide variety of cell types, including airway epithelium (Rosenfeld *et al.* (1992) cited *supra*), endothelial cells (Lemarchand *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:6482-6486), hepatocytes (Herz and Gerard (1993) *Proc. Natl. Acad. Sci. USA* 90:2812-2816) and muscle cells (Quantin *et al.*
- 35 (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584). Additionally, introduced adenoviral DNA (and foreign DNA contained therein) is not integrated into the genome of a host cell but remains episomal, thereby avoiding potential problems that can occur as a result of insertional mutagenesis in situations where introduced DNA becomes integrated into the host genome (*e.g.*, retroviral DNA). Moreover, the carrying capacity of the adenoviral genome for

foreign DNA is large (up to 8 kilobases) relative to other gene delivery vectors (Berkner *et al.* cited *supra*; Haj-Ahmand and Graham (1986) *J. Virol.* 57:267). Most replication-defective adenoviral vectors currently in use are deleted for all or parts of the viral E1 and E3 genes but retain as much as 80 % of the adenoviral genetic material.

- 5        *Adeno-Associated Viruses*: Adeno-associated virus (AAV) is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle. (For a review see Muzyczka *et al.* *Curr. Topics in Micro. and Immunol.* (1992) 158:97-129). It is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable
- 10 integration (see for example Flotte *et al.* (1992) *Am. J. Respir. Cell. Mol. Biol.* 7:349-356; Samulski *et al.* (1989) *J. Virol.* 63:3822-3828; and McLaughlin *et al.* (1989) *J. Virol.* 62:1963-1973). Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate. Space for exogenous DNA is limited to about 4.5 kb. An AAV vector such as that described in Tratschin *et al.* (1985) *Mol. Cell. Biol.* 5:3251-3260 can be used to introduce
- 15 DNA into cells. A variety of nucleic acids have been introduced into different cell types using AAV vectors (see for example Hermonat *et al.* (1984) *Proc. Natl. Acad. Sci. USA* 81:6466-6470; Tratschin *et al.* (1985) *Mol. Cell. Biol.* 4:2072-2081; Wondisford *et al.* (1988) *Mol. Endocrinol.* 2:32-39; Tratschin *et al.* (1984) *J. Virol.* 51:611-619; and Flotte *et al.* (1993) *J. Biol. Chem.* 268:3781-3790).
- 20        The efficacy of a particular expression vector system and method of introducing nucleic acid into a cell can be assessed by standard approaches routinely used in the art. For example, DNA introduced into a cell can be detected by a filter hybridization technique (*e.g.*, Southern blotting) and RNA produced by transcription of introduced DNA can be detected, for example, by Northern blotting, RNase protection or reverse transcriptase-polymerase
- 25 chain reaction (RT-PCR). The gene product can be detected by an appropriate assay, for example by immunological detection of a produced protein, such as with a specific antibody, or by a functional assay to detect a functional activity of the gene product, such as an enzymatic assay.

- 30        A modulatory agent, such as a chemical compound that stimulates or inhibits endogenous c-Maf activity, can be administered to a subject as a pharmaceutical composition. Such compositions typically comprise the modulatory agent and a pharmaceutically acceptable carrier. As used herein the term "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical
- 35 administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. For example, solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

### III. Applications of the Methods of the Invention

Identification of the transcription factor that controls the production of IL-4, and hence continued formation of Th2 cells, allows for selective manipulation of T cell subsets in a variety of clinical situations using the modulatory methods of the invention. The stimulatory methods of the invention (*i.e.*, methods that use a stimulatory agent) result in

production of Th2-associated cytokines, with concomitant promotion of a Th2 response and downregulation of a Th1 response. In contrast, the inhibitory methods of the invention (*i.e.*, methods that use an inhibitory agent) inhibit the production of Th2-associated cytokines, with concomitant downregulation of a Th2 response and promotion of a Th1 response. Thus, to  
5 treat a disease condition wherein a Th2 response is beneficial, a stimulatory method of the invention is selected such that Th2 responses are promoted while downregulating Th1 responses. Alternatively, to treat a disease condition wherein a Th1 response is beneficial, an inhibitory method of the invention is selected such that Th2 responses are downregulated while promoting Th1 responses. Application of the methods of the invention to the treatment  
10 of disease conditions may result in cure of the condition, a decrease in the type or number of symptoms associated with the condition, either in the long term or short term (*i.e.*, amelioration of the condition) or simply a transient beneficial effect to the subject.

Numerous disease conditions associated with a predominant Th1 or Th2-type response have been identified and could benefit from modulation of the type of response  
15 mounted in the individual suffering from the disease condition. Application of the immunomodulatory methods of the invention to such diseases is described in further detail below.

#### A. Allergies

20 Allergies are mediated through IgE antibodies whose production is regulated by the activity of Th2 cells and the cytokines produced thereby. In allergic reactions, IL-4 is produced by Th2 cells, which further stimulates production of IgE antibodies and activation of cells that mediate allergic reactions, *i.e.*, mast cells and basophils. IL-4 also plays an  
25 important role in eosinophil mediated inflammatory reactions. Accordingly, the inhibitory methods of the invention can be used to inhibit the production of Th2-associated cytokines, and in particular IL-4, in allergic patients as a means to downregulate production of pathogenic IgE antibodies. An inhibitory agent may be directly administered to the subject or cells (*e.g.*, Thp cells or Th2 cells) may be obtained from the subject, contacted with an  
30 inhibitory agent *ex vivo*, and readministered to the subject. Moreover, in certain situations it may be beneficial to coadminister to the subject the allergen together with the inhibitory agent or cells treated with the inhibitory agent to inhibit (*e.g.*, desensitize) the allergen-specific response. The treatment may be further enhanced by administering other Th1-promoting agents, such as the cytokine IL-12 or antibodies to Th2-associated cytokines (*e.g.*,  
35 anti-IL-4 antibodies), to the allergic subject in amounts sufficient to further stimulate a Th1-type response.

## B. Cancer

The expression of Th2-promoting cytokines has been reported to be elevated in cancer patients (see *e.g.*, Yamamura, M., *et al.* (1993) *J. Clin. Invest.* 91:1005-1010; Pisa, P., *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:7708-7712) and malignant disease is often associated with a shift from Th1 type responses to Th2 type responses along with a worsening of the course of the disease. Accordingly, the inhibitory methods of the invention can be used to inhibit the production of Th2-associated cytokines in cancer patients, as a means to counteract the Th1 to Th2 shift and thereby promote an ongoing Th1 response in the patients to ameliorate the course of the disease. The inhibitory method can involve either direct administration of an inhibitory agent to a subject with cancer or *ex vivo* treatment of cells obtained from the subject (*e.g.*, Thp or Th2 cells) with an inhibitory agent followed by readministration of the cells to the subject. The treatment may be further enhanced by administering other Th1-promoting agents, such as the cytokine IL-12 or antibodies to Th2-associated cytokines (*e.g.*, anti-IL-4 antibodies), to the recipient in amounts sufficient to further stimulate a Th1-type response.

## C. Infectious Diseases

The expression of Th2-promoting cytokines also has been reported to increase during a variety of infectious diseases, including HIV infection, tuberculosis, leishmaniasis, schistosomiasis, filarial nematode infection and intestinal nematode infection (see *e.g.*; Shearer, G.M. and Clerici, M. (1992) *Prog. Chem. Immunol.* 54:21-43; Clerici, M and Shearer, G.M. (1993) *Immunology Today* 14:107-111; Fauci, A.S. (1988) *Science* 239:617-623; Locksley, R. M. and Scott, P. (1992) *Immunoparasitology Today* 1:A58-A61; Pearce, E.J., *et al.* (1991) *J. Exp. Med.* 173:159-166; Grzych, J-M., *et al.* (1991) *J. Immunol.* 141:1322-1327; Kullberg, M.C., *et al.* (1992) *J. Immunol.* 148:3264-3270; Bancroft, A.J., *et al.* (1993) *J. Immunol.* 150:1395-1402; Pearlman, E., *et al.* (1993) *Infect. Immun.* 61:1105-1112; Else, K.J., *et al.* (1994) *J. Exp. Med.* 179:347-351) and such infectious diseases are also associated with a Th1 to Th2 shift in the immune response. Accordingly, the inhibitory methods of the invention can be used to inhibit the production of Th2-associated cytokines in subjects with infectious diseases, as a means to counteract the Th1 to Th2 shift and thereby promote an ongoing Th1 response in the patients to ameliorate the course of the infection. The inhibitory method can involve either direct administration of an inhibitory agent to a subject with an infectious disease or *ex vivo* treatment of cells obtained from the subject (*e.g.*, Thp or Th2 cells) with an inhibitory agent followed by readministration of the cells to the subject. The treatment may be further enhanced by administering other Th1-promoting agents, such as the cytokine IL-12 or antibodies to Th2-associated cytokines (*e.g.*, anti-IL-4 antibodies), to the recipient in amounts sufficient to further stimulate a Th1-type response.

#### D. Autoimmune Diseases

The stimulatory methods of the invention can be used therapeutically in the treatment of autoimmune diseases that are associated with a Th2-type dysfunction. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and that promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Modulation of T helper-type responses can have an effect on the course of the autoimmune disease. For example, in experimental allergic encephalomyelitis (EAE), stimulation of a Th2-type response by administration of IL-4 at the time of the induction of the disease diminishes the intensity of the autoimmune disease (Paul, W.E., *et al.* (1994) *Cell* 76:241-251). Furthermore, recovery of the animals from the disease has been shown to be associated with an increase in a Th2-type response as evidenced by an increase of Th2-specific cytokines (Koury, S. J., *et al.* (1992) *J. Exp. Med.* 176:1355-1364). Moreover, T cells that can suppress EAE secrete Th2-specific cytokines (Chen, C., *et al.* (1994) *Immunity* 1:147-154). Since stimulation of a Th2-type response in EAE has a protective effect against the disease, stimulation of a Th2 response in subjects with multiple sclerosis (for which EAE is a model) is likely to be beneficial therapeutically.

Similarly, stimulation of a Th2-type response in type I diabetes in mice provides a protective effect against the disease. Indeed, treatment of NOD mice with IL-4 (which promotes a Th2 response) prevents or delays onset of type I diabetes that normally develops in these mice (Rapoport, M.J., *et al.* (1993) *J. Exp. Med.* 178:87-99). Thus, stimulation of a Th2 response in a subject suffering from or susceptible to diabetes may ameliorate the effects of the disease or inhibit the onset of the disease.

Yet another autoimmune disease in which stimulation of a Th2-type response may be beneficial is rheumatoid arthritis (RA). Studies have shown that patients with rheumatoid arthritis have predominantly Th1 cells in synovial tissue (Simon, A.K., *et al.*, (1994) *Proc. Natl. Acad. Sci. USA* 91:8562-8566). By stimulating a Th2 response in a subject with RA, the detrimental Th1 response can be concomitantly downmodulated to thereby ameliorate the effects of the disease.

Accordingly, the stimulatory methods of the invention can be used to stimulate production of Th2-associated cytokines in subjects suffering from, or susceptible to, an autoimmune disease in which a Th2-type response is beneficial to the course of the disease. The stimulatory method can involve either direct administration of a stimulatory agent to the subject or *ex vivo* treatment of cells obtained from the subject (*e.g.*, Thp, Th1 cells, B cells, non-lymphoid cells) with a stimulatory agent followed by readministration of the cells to the subject. The treatment may be further enhanced by administering other Th2-promoting agents, such as IL-4 itself or antibodies to Th1-associated cytokines, to the subject in amounts sufficient to further stimulate a Th2-type response.

In contrast to the autoimmune diseases described above in which a Th2 response is desirable, other autoimmune diseases may be ameliorated by a Th1-type response. Such diseases can be treated using an inhibitory agent of the invention (as described above for cancer and infectious diseases). The treatment may be further enhanced by administering a Th1-promoting cytokine (*e.g.*, IFN- $\gamma$ ) to the subject in amounts sufficient to further stimulate a Th1-type response.

The efficacy of agents for treating autoimmune diseases can be tested in the above described animal models of human diseases (*e.g.*, EAE as a model of multiple sclerosis and the NOD mice as a model for diabetes) or other well characterized animal models of human autoimmune diseases. Such animal models include the *mrl/lpr/lpr* mouse as a model for lupus erythematosus, murine collagen-induced arthritis as a model for rheumatoid arthritis, and murine experimental myasthenia gravis (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 840-856). A modulatory (*i.e.*, stimulatory or inhibitory) agent of the invention is administered to test animals and the course of the disease in the test animals is then monitored by the standard methods for the particular model being used. Effectiveness of the modulatory agent is evidenced by amelioration of the disease condition in animals treated with the agent as compared to untreated animals (or animals treated with a control agent).

Non-limiting examples of autoimmune diseases and disorders having an autoimmune component that may be treated according to the invention include diabetes mellitus, arthritis (including rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, psoriatic arthritis), multiple sclerosis, myasthenia gravis, systemic lupus erythematosus, autoimmune thyroiditis, dermatitis (including atopic dermatitis and eczematous dermatitis), psoriasis, Sjögren's Syndrome, including keratoconjunctivitis sicca secondary to Sjögren's Syndrome, alopecia areata, allergic responses due to arthropod bite reactions, Crohn's disease, aphthous ulcer, iritis, conjunctivitis, keratoconjunctivitis, ulcerative colitis, asthma, allergic asthma, cutaneous lupus erythematosus, scleroderma, vaginitis, proctitis, drug eruptions, leprosy reversal reactions, erythema nodosum leprosum, autoimmune uveitis, allergic encephalomyelitis, acute necrotizing hemorrhagic encephalopathy, idiopathic bilateral progressive sensorineural hearing loss, aplastic anemia, pure red cell anemia, idiopathic thrombocytopenia, polychondritis, Wegener's granulomatosis, chronic active hepatitis, Stevens-Johnson syndrome, idiopathic sprue, lichen planus, Crohn's disease, Graves ophthalmopathy, sarcoidosis, primary biliary cirrhosis, uveitis posterior, and interstitial lung fibrosis.

#### E. Transplantation

While graft rejection or graft acceptance may not be attributable exclusively to the action of a particular T cell subset (*i.e.*, Th1 or Th2 cells) in the graft recipient (for a



discussion see Dallman, M.J. (1995) *Curr. Opin. Immunol.* 7:632-638), numerous studies have implicated a predominant Th2 response in prolonged graft survival or a predominant Th2 response in graft rejection. For example, graft acceptance has been associated with production of a Th2 cytokine pattern and/or graft rejection has been associated with production of a Th1 cytokine pattern (see *e.g.*, Takeuchi, T. *et al.* (1992) *Transplantation* 53:1281-1291; Tzakis, A.G. *et al.* (1994) *J. Pediatr. Surg.* 29:754-756; Thai, N.L. *et al.* (1995) *Transplantation* 59:274-281). Additionally, adoptive transfer of cells having a Th2 cytokine phenotype prolongs skin graft survival (Maeda, H. *et al.* (1994) *Int. Immunol.* 6:855-862) and reduces graft-versus-host disease (Fowler, D.H. *et al.* (1994) *Blood* 84:3540-3549; Fowler, D.H. *et al.* (1994) *Prog. Clin. Biol. Res.* 389:533-540). Still further, administration of IL-4, which promotes Th2 differentiation, prolongs cardiac allograft survival (Levy, A.E. and Alexander, J.W. (1995) *Transplantation* 60:405-406), whereas administration of IL-12 in combination with anti-IL-10 antibodies, which promotes Th1 differentiation, enhances skin allograft rejection (Gorczynski, R.M. *et al.* (1995) *Transplantation* 60:1337-1341).

Accordingly, the stimulatory methods of the invention can be used to stimulate production of Th2-associated cytokines in transplant recipients to prolong survival of the graft. The stimulatory methods can be used both in solid organ transplantation and in bone marrow transplantation (*e.g.*, to inhibit graft-versus-host disease). The stimulatory method can involve either direct administration of a stimulatory agent to the transplant recipient or *ex vivo* treatment of cells obtained from the subject (*e.g.*, Thp, Th1 cells, B cells, non-lymphoid cells) with a stimulatory agent followed by readministration of the cells to the subject. The treatment may be further enhanced by administering other Th2-promoting agents, such as IL-4 itself or antibodies to Th1-associated cytokines, to the recipient in amounts sufficient to further stimulate a Th2-type response.

In addition to the foregoing disease situations, the modulatory methods of the invention also are useful for other purposes. For example, the stimulatory methods of the invention (*i.e.*, methods using a stimulatory agent) can be used to stimulate production of Th2-promoting cytokines (*e.g.*, IL-4) *in vitro* for commercial production of these cytokines (*e.g.*, cells can be contacted with the stimulatory agent *in vitro* to stimulate IL-4 production and the IL-4 can be recovered from the culture supernatant, further purified if necessary, and packaged for commercial use).

Furthermore, the modulatory methods of the invention can be applied to vaccinations to promote either a Th1 or a Th2 response to an antigen of interest in a subject. That is, the agents of the invention can serve as adjuvants to direct an immune response to a vaccine either to a Th1 response or a Th2 response. For example, to stimulate an antibody response to an antigen of interest (*i.e.*, for vaccination purposes), the antigen and a stimulatory agent of the invention can be coadministered to a subject to promote a Th2 response to the antigen in the subject, since Th2 responses provide efficient B cell help and promote IgG1 production.

Alternatively, to promote a cellular immune response to an antigen of interest, the antigen and an inhibitory agent of the invention can be coadministered to a subject to promote a Th1 response to the antigen in a subject, since Th1 responses favor the development of cell-mediated immune responses (*e.g.*, delayed hypersensitivity responses). The antigen of interest and the modulatory agent can be formulated together into a single pharmaceutical composition or in separate compositions. In a preferred embodiment, the antigen of interest and the modulatory agent are administered simultaneously to the subject. Alternatively, in certain situations it may be desirable to administer the antigen first and then the modulatory agent or vice versa (for example, in the case of an antigen that naturally evokes a Th1 response, it may be beneficial to first administer the antigen alone to stimulate a Th1 response and then administer a stimulatory agent, alone or together with a boost of antigen, to shift the immune response to a Th2 response).

#### IV. Compositions for Modulating Th2-Associated Cytokine Production

Another aspect of the invention pertains to compositions that can be used to modulate Th2-associated cytokine production by a cell or Th1/Th2 development in a subject in accordance with the methods of the invention. The invention provides recombinant expression vectors comprising a nucleotide sequence encoding a maf family protein operatively linked to regulatory sequences that direct expression of the maf family protein specifically in certain cell types. In a preferred embodiment, the regulatory sequences direct expression of the maf family protein specifically in lymphoid cells (*e.g.*, T cells or B cells). In one embodiment, the lymphoid cells are T cells. T cell specific regulatory elements are known in the art, such as the promoter regulatory region of T cell receptor genes (see *e.g.*, Winoto and Baltimore (1989) *EMBO J.* 8:729-733; Leiden, J.M. (1994) *Annu. Rev. Immunol.* 11:539-570; Hettman, T. and Cohen, A. (1994) *Mol. Immunol.* 31:315-322; Redondo, J.M. *et al.* (1991) *Mol. Cell. Biol.* 11:5671-5680). Other examples of T cell specific regulatory elements are those derived from the CD3 gene (see *e.g.*, Clevers, H. *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:8623-8627; Clevers, H.C. *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:8156-8160; Georgopoulos, K. *et al.* (1988) *EMBO J.* 7:2401-2407), the CD4 gene (see *e.g.*, Sawada, S. and Littman, D.R. (1991) *Mol. Cell. Biol.* 11:5506-5515; Salmon, P. *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:7739-7743; Hanna, Z. *et al.* (1994) *Mol. Cell. Biol.* 14:1084-1094; see also GenBank accession numbers U01066 and S68043 for human CD4 promoter sequences) and the CD2 gene (see *e.g.*, Zhumabekov, T. *et al.* (1995) *J. Immunol. Methods* 185:133-140). A DNA fragment comprising one or more T cell specific regulatory elements, such as a promoter and enhancer of a T cell receptor gene, can be obtained by standard molecular biology methods, such as by PCR using oligonucleotide primers corresponding to the 5' and 3' ends of the desired region and genomic DNA from T cells as the template. Once the DNA fragment comprising T cell specific regulatory elements is obtained, it can be operatively linked to a cDNA encoding a maf protein (*e.g.*, the two DNA

fragments can be ligated together such that the regulatory elements are located 5' of the maf sequences) and introduced into vector, such as a plasmid vector, using standard molecular biology techniques.

5 In another embodiment, the lymphoid cells are B cells (i.e., within the recombinant expression vector the nucleotide sequences encoding a maf family protein are operatively linked to regulatory sequences that direct expression of the maf family specifically in B cells). B cell specific regulatory elements are known in the art, such as the promoter regulatory region of immunoglobulin genes (see *e.g.*, Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748). Other examples of B cell specific regulatory elements are those derived from the CD20 (B1) gene (see *e.g.*, Thevenin, C. *et al.* (1993) *J. Biol. Chem.* 268:5949-5956; Rieckmann, P. *et al.* (1991) *J. Immunol.* 147:3994-3999), the Fc epsilon RIIa gene (see *e.g.*, Suter, U. *et al.* (1989) *J. Immunol.* 143:3087-3092) and major histocompatibility class II genes (see *e.g.*, Glimcher, L.H. and Kara, C.J. (1992) *Annu. Rev. Immunol.* 10:13-49; Benoist, C. and Mathis, D. (1990) *Annu. Rev. Immunol.* 8:681-715). A DNA fragment comprising B cell specific regulatory elements, such as a promoter and enhancer of an immunoglobulin gene, can be obtained by standard molecular biology methods, such as by PCR using oligonucleotide primers corresponding to the 5' and 3' ends of the desired region and genomic DNA from B cells as the template. Once the DNA fragment comprising B cell specific regulatory elements is obtained, it can be operatively linked to a cDNA encoding a maf protein (*e.g.*, the two DNA fragments can be ligated together such that the regulatory elements are located 5' of the maf sequences) and introduced into vector, such as a plasmid vector, using standard molecular biology techniques.

25 In yet another embodiment, the invention provides recombinant expression vectors comprising a nucleotide sequence encoding a maf family protein operatively linked to regulatory sequences that direct expression of the maf family protein specifically in hematopoietic stem cells. Hematopoietic stem cell specific regulatory elements are known in the art. Preferably regulatory elements derived from the CD34 gene are used (see *e.g.*, Satterthwaite, A.B. *et al.* (1992) *Genomics* 12:788-794; Burn, T. C. *et al.* (1992) *Blood* 80:3051-3059).

30 Another aspect of the invention pertains to recombinant host cells that express a maf family protein. Such host cells can be used to produce a Th2-associated cytokine (*e.g.*, IL-4). Such host cells also can be administered to a subject to produce a Th2-associated cytokine in the subject as a means to manipulate Th1:Th2 ratios in the subject. The terms "host cell" and "recombinant host cell" are used interchangeably herein to refer to a cell into which a recombinant expression vector has been introduced. It is understood that such terms refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but as long as these

progeny cells retain the recombinant expression vector, these progeny are still intended to be included within the scope of the term "host cell" as used herein.

In one embodiment, the invention provides a host lymphoid cell into which a recombinant expression vector encoding a maf family protein has been introduced. The host lymphoid cell can be a T cell or a B cell. A host T cell of the invention can be, for example a T cell clone that is cultured *in vitro* (such as those described in the Examples) or, alternatively, a normal T cell that is isolated from a subject (*e.g.*, a peripheral blood T cell or a splenic T cell). Standard methods for preparing and culturing T cell clones *in vitro*, or isolating T cells (*e.g.*, from peripheral blood) are known in the art, for example through the use of mAbs that bind T cell specific cell surface markers (*e.g.*, CD3) or surface markers for specific subsets of T cells (*e.g.*, CD4 or CD8). The recombinant expression vector can be introduced into the T cell by one of a variety of known transfection methods for introducing DNA into mammalian cells, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory manuals.

In another embodiment, the host lymphoid cell of the invention is a host B cell into which a recombinant expression vector encoding a maf family protein has been introduced. The B cell can be, for example a B lymphoma cell that is cultured *in vitro* (such as M12 cells as described in the Examples) or, alternatively, a normal B cell that is isolated from a subject (*e.g.*, a peripheral blood B cell or a splenic B cell). Various B lymphoma cell lines are available in the art and standard methods for culturing such cells *in vitro* are known. Additionally, standard methods for isolating normal B cells (*e.g.*, from peripheral blood) are known in the art, for example through the use of mAbs that bind B cell specific cell surface markers (*e.g.*, membrane immunoglobulin, B7-1, CD20). The recombinant expression vector can be introduced into the B cell by standard methods, as described above for T cells.

In yet another embodiment, the invention provides a host hematopoietic stem cell into which a recombinant expression vector encoding a maf family protein has been introduced. Hematopoietic stem cells can be isolated from a subject (*e.g.*, from peripheral blood or bone marrow of the subject) using standard methods known in the art for isolating such stem cells, for example through the use of mAbs that bind hematopoietic stem cell specific cell surface markers, preferably CD34 (for further descriptions of isolation of stem cells, see *e.g.*, Wagner, J.E. *et al.* (1995) *Blood* 86:512-523; Murray, L. *et al.* (1995) *Blood* 85:368-378; Bernardi, A.C. *et al.* (1995) *Science* 267:104-108; Bernstein, I.D. *et al.* (1994) *Blood Cells* 20:15-24; Angelini, A. *et al.* (1993) *Int. J. Artif. Organs* 16 Suppl. 5:13-18; Kato, K. and Radburch, A. (1993) *Cytometry* 14:384-392; Lebkowski, J.S. *et al.* (1992) *Transplantation* 53:1011-1019; Lebkowski, J. *et al.* (1993) *J. Hematother.* 2:339-342). The recombinant

expression vector can be introduced into the hematopoietic stem cell by standard methods, as described above for T cells.

#### V. Screening Assays

5        Another aspect of the invention pertains to screening assays for identifying compounds that modulate the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene. In various embodiments, these screening assays can identify, for example, compounds that modulate the expression or functional activity of the transcription factor, proteins that interact with the transcription factor, as well as compounds  
10        that modulate these protein-protein interactions, and compounds that modulate the interaction of the transcription factor with a MARE within a Th2-associated cytokine gene.

      In a preferred embodiment, the invention provides a method comprising:

- a) preparing an indicator cell, wherein said indicator cell contains:
  - 15        i) a recombinant expression vector encoding a transcription factor that regulates expression of a Th2-associated cytokine gene; and
  - ii) a vector comprising regulatory sequences of the Th2-associated cytokine gene operatively linked a reporter gene;
- b) contacting the indicator cell with a test compound;
- c) determining the level of expression of the reporter gene in the indicator cell in the  
20        presence of the test compound;
- d) comparing the level of expression of the reporter gene in the indicator cell in the presence of the test compound with the level of expression of the reporter gene in the indicator cell in the absence of the test compound; and
- e) identifying a compound that modulates the activity of a transcription factor that  
25        regulates expression of a Th2-associated cytokine gene.

      Preferably, the transcription factor is a member of the maf family and most preferably a c-Maf protein. Recombinant expression vectors that can be used for expression of a c-Maf protein in an indicator cell are known in the art (see discussions above and also the  
30        Examples). In one embodiment, within the expression vector the c-Maf coding sequences are operatively linked to regulatory sequences that allow for constitutive expression of c-Maf in the indicator cell (*e.g.*, viral regulatory sequences, such as a cytomegalovirus promoter/enhancer, can be used). Use of a recombinant expression vector that allows for constitutive expression of c-Maf in the indicator cell is preferred for identification of  
35        compounds that enhance or inhibit the activity of c-Maf. In an alternative embodiment, within the expression vector the c-Maf coding sequences are operatively linked to regulatory sequences of the *c-maf* gene (*i.e.*, the promoter regulatory region derived from the endogenous *c-maf* gene). Use of a recombinant expression vector in which c-Maf protein

expression is controlled by *c-maf* regulatory sequences is preferred for identification of compounds that enhance or inhibit the transcriptional expression of c-Maf.

Preferably, the Th2-associated cytokine is interleukin-4. It has previously shown that Th2-specific, inducible IL-4 expression can be directed by as little as 157 bp of the proximal IL-4 promoter in Th2 cells (Hodge, M. *et al.* (1995) *J. Immunol.* 154:6397-6405).

Accordingly, in one embodiment, the method utilizes a reporter gene construct containing this region of the proximal IL-4 promoter, most preferably nucleotides -157 to +58 (relative to the start site of transcription at +1) of the IL-4 promoter. Alternatively, stronger reporter gene expression can be achieved using a longer portion of the IL-4 upstream regulatory region, such as about 3 kb of upstream regulatory sequences. Suitable reporter gene constructs are described in Todd, M. *et al.* (1993) *J. Exp. Med.* 177:1663-1674. See also the Examples for descriptions of IL-4 reporter gene constructs.

A variety of reporter genes are known in the art and are suitable for use in the screening assays of the invention. Examples of suitable reporter genes include those which encode chloramphenicol acetyltransferase, beta-galactosidase, alkaline phosphatase or luciferase. Standard methods for measuring the activity of these gene products are known in the art.

A variety of cell types are suitable for use as an indicator cell in the screening assay. Preferably a cell line is used which does not normally express c-Maf, such as a B cell (*e.g.*, the M12 B lymphoma cell line) or a Th1 cell clone (*e.g.*, AE7 cells). Nonlymphoid cell lines can also be used as indicator cells, such as the HepG2 hepatoma cell line.

In one embodiment, the level of expression of the reporter gene in the indicator cell in the presence of the test compound is higher than the level of expression of the reporter gene in the indicator cell in the absence of the test compound and the test compound is identified as a compound that stimulates the expression or activity of the transcription factor. In another embodiment, the level of expression of the reporter gene in the indicator cell in the presence of the test compound is lower than the level of expression of the reporter gene in the indicator cell in the absence of the test compound and the test compound is identified as a compound that inhibits the expression or activity of the transcription factor.

Alternative to the use of a reporter gene construct, compounds that modulate the expression or activity of c-Maf can be identified by using other "read-outs." For example, an indicator cell can be transfected with a c-Maf expression vector, incubated in the presence and in the absence of a test compound, and Th2-associated cytokine production can be assessed by detecting cytokine mRNA (*e.g.*, IL-4 mRNA) in the indicator cell or cytokine secretion (*i.e.*, IL-4 secretion) into the culture supernatant. Standard methods for detecting cytokine mRNA, such as reverse transcription-polymerase chain reaction (RT-PCR) are known in the art. Standard methods for detecting cytokine protein in culture supernatants, such as enzyme linked immunosorbent assays (ELISA) are also known in the art. For further descriptions of methods for detecting cytokine mRNA and/or protein, see also the Examples.

In another embodiment, the invention provides a screening assay for identifying proteins in Th2 cells that interact with c-Maf. These assays can be designed based on the two-hybrid assay system (also referred to as an interaction trap assay) known in the art (see *e.g.*, Field U.S. Patent No. 5,283,173; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; and Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696). The two-hybrid assay is generally used for identifying proteins that interact with a particular target protein. The assay employs gene fusions to identify proteins capable of interacting to reconstitute a functional transcriptional activator. The transcriptional activator consists of a DNA-binding domain and a transcriptional activation domain, wherein both domains are required to activate transcription of genes downstream from a target sequence (such as an upstream activator sequence (UAS) for GAL4). DNA sequences encoding a target "bait" protein are fused to either of these domains and a library of DNA sequences is fused to the other domain. "Fish" fusion proteins (generated from the fusion library) capable of binding to the target-fusion protein (*e.g.*, a target GAL4-fusion "bait") will generally bring the two domains (DNA-binding domain and transcriptional activation domain) into close enough proximity to activate the transcription of a reporter gene inserted downstream from the target sequence. Thus, the "fish" proteins can be identified by their ability to reconstitute a functional transcriptional activator (*e.g.*, a functional GAL4 transactivator).

This general two-hybrid system can be applied to the identification of proteins in Th2 cells that interact with c-Maf by construction of a target c-Maf fusion protein (*e.g.*, a c-Maf/GAL4 binding domain fusion as the "bait") and a cDNA library of "fish" fusion proteins (*e.g.*, a cDNA/GAL4 activation domain library), wherein the cDNA library is prepared from mRNA of Th2 cells, and introducing these constructs into a host cell that also contains a reporter gene construct linked to a regulatory sequence responsive to c-Maf (*e.g.*, a MARE sequence, for example a region of the IL-4 promoter, as discussed above). cDNAs encoding proteins from Th2 cells that interact with c-Maf can be identified based upon transactivation of the reporter gene construct. Accordingly, the invention provides a method for identifying a protein in a Th2 cell that interacts with c-Maf comprising:

- a) providing a two hybrid assay including a host cell which contains
  - i) a reporter gene operably linked to a transcriptional regulatory sequence;
  - ii) a first chimeric gene which encodes a first fusion protein, said first fusion protein including c-Maf;
  - iii) a library of second chimeric genes which encodes second fusion proteins, the second fusion proteins including proteins derived from Th2 cells;

wherein expression of the reporter gene is sensitive to interactions between the first fusion protein, the second fusion protein and the transcriptional regulatory sequence;

- b) determining the level of expression of the reporter gene in the host cell; and
- c) identifying a protein in a Th2 cell that interacts with c-Maf.

Alternatively, a "single-hybrid" assay, such as that described in Sieweke, M.H. *et al.* (1996) *Cell* 85:49-60, can be used to identify proteins from Th2 cells that interact with c-Maf. This assay is a modification of the two-hybrid system discussed above. In this system, the "bait" is a transcription factor from which the transactivation domain has been removed (*e.g.*, c-Maf from which the amino-terminal transactivation domain has been removed) and the "fish" is a non-fusion cDNA library (*e.g.*, a cDNA library prepared from Th2 cells). These constructs are introduced into host cells (*e.g.*, yeast cells) that also contains a reporter gene construct linked to a regulatory sequence responsive to the transcription factor (*e.g.*, a MARE sequence, for example a region of the IL-4 promoter, responsive to c-Maf). cDNAs encoding proteins from Th2 cells that interact with c-Maf can be identified based upon transactivation of the reporter gene construct.

In yet another embodiment, the invention provides a screening assay for identifying compounds that modulate the interaction of c-Maf with a MARE in an IL-4 gene regulatory region. Assays are known in the art that detect the interaction of a DNA binding protein with a target DNA sequence (*e.g.*, electrophoretic mobility shift assays, DNase I footprinting assays and the like; for further descriptions see the Examples). By performing such assays in the presence and absence of test compounds, these assays can be used to identify compounds that modulate (*e.g.*, inhibit or enhance) the interaction of the DNA binding protein with its target DNA sequence. Accordingly, the invention provides a method for identifying a compound that modulates the interaction of a c-Maf protein with a maf response element (MARE) of an IL-4 gene regulatory region, comprising:

- a) providing a c-Maf protein and a DNA fragment comprising a MARE of an IL-4 gene regulatory region;
- b) incubating the c-Maf protein and DNA fragment in the presence of a test compound;
- c) determining the amount of binding of the c-Maf protein to the DNA fragment in the presence of the test compound;
- d) comparing the amount of binding of the c-Maf protein to the DNA fragment in the presence of the test compound with the amount of binding of the c-Maf protein to the DNA fragment in the absence of the test compound; and
- e) identifying a compound that modulates the interaction of a c-Maf protein with a MARE of an IL-4 gene regulatory region.

In one embodiment, the amount of binding of the c-Maf protein to the DNA fragment in the presence of the test compound is greater than the amount of binding of the c-Maf protein to the DNA fragment in the absence of the test compound, in which case the test compound is identified as a compound that enhances binding of c-Maf to the MARE. In



another embodiment, the amount of binding of the c-Maf protein to the DNA fragment in the presence of the test compound is less than the amount of binding of the c-Maf protein to the DNA fragment in the absence of the test compound, in which case the test compound is identified as a compound that inhibits binding of c-Maf to the MARE.

5

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference. Nucleotide and amino acid sequences deposited in public databases as referred to herein are also hereby incorporated by reference.

10

**EXAMPLE 1: Cytokine Specificity is Due to a Positive Transacting Factor and Not to a Repressor**

15 Tissue specificity can be achieved through the action of repressor or silencer proteins. Thus it was possible that the IL-2 and IL-4 genes were actively repressed in Th2 and Th1 cells respectively. To test for the existence of repressor proteins, somatic cell fusions were performed between a Th1(D1.1) and a Th2 (D10) clone of differing MHC Class I haplotypes. The Th1 clone D1.1 (K<sup>d</sup>) and the Th2 clone D10 (K<sup>k</sup>) were fused according to the  
20 "suspension cell fusion" procedure (Lane, R.D. *et al.* (1986) *Methods Enzymol.* 121:183-192). After fusion, the cells were allowed to recover for 8 hours and then double- stained using PE-conjugated anti-K<sup>k</sup> and FITC-conjugated anti-K<sup>d</sup> antibodies (Pharmingen, La Jolla, CA). Cells were then sorted on the basis of size to distinguish unfused cells from hetero and homokaryons and by fluorescence to identify single-positive and double-positive cells. As  
25 indicated in the schematic of this approach shown in Figure 1A, three populations were sorted for: large PE-positive cells (D1.1 x D1.1), large FITC-positive cells (D10 x D10), and large PE and FITC positive cells (D1.1 x D10). Cells expressing both MHC class I K<sup>b</sup> and K<sup>k</sup> markers were heterokaryons while cells expressing only K<sup>b</sup> or K<sup>k</sup> represented homokaryons and served as controls.

30 The three populations were then stimulated in culture with antibodies to CD3 to activate cytokine gene expression and RNA prepared for RT-PCR and Northern blot analysis. Approximately  $5 \times 10^5$  cells were obtained for each population. Routinely, 5-10% of the cells had undergone fusion. Each of these three populations was then split in half, one half transferred to pre-rinsed anti-CD3 coated plates, the remaining half to uncoated plates. After  
35 four hours, the cells were harvested, and poly(A<sup>+</sup>) RNA isolated using the Micro-FastTrackTM kit (Stratagene, La Jolla, CA). cDNA was made using the SuperScript kit (Gibco/BRL, Bethesda, MD), and used for PCR analysis using commercially available primers specific for murine IL-2, IL-4 and  $\beta$ -actin according to the manufacturer's instructions (Stratagene, La Jolla, CA). PCR reactions included 0.5  $\mu$ Ci  $\alpha$ -<sup>32</sup>P-dCTP (3000

662720" 9948486

Ci/mmol, NEN Dupont). PCR products were ethanol precipitated, separated by nondenaturing PAGE and dried and visualized by autoradiography.

The results of the RT-PCT analysis of cytokine mRNA expression are shown in Figure 1B. The Th1 and Th2 clones and the Th homokaryons transcribed only IL-2 (Th1) or IL-4 (Th2) respectively, while the Th1/Th2 heterokaryons produced both cytokines. In contrast, the existence of repressor protein(s) should have resulted in the extinction of both cytokines in the heterokaryons. From these experiments, it was concluded that cytokine specificity in Th1 vs. Th2 cells was mediated by Th-specific positive transacting factors rather than by selective silencer proteins.

#### **EXAMPLE 2: Isolation of a Th2-Specific *c-maf* Gene from a cDNA Library Prepared from an Anti-CD3 Activated Th2 Clone**

In the course of screening a cDNA library prepared from an anti-CD3 activated Th2 clone, D10, for NF-AT-interacting proteins by the yeast two-hybrid system (for descriptions of this system, see *e.g.*, Field U.S. Patent No. 5,283,173; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; and Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696), multiple cDNAs were isolated, all of which were extremely weak interactors. All cDNAs obtained in this screen were next evaluated for Th-specific expression by Northern blot analysis using a panel of Th1 and Th2 clones. One such cDNA, which was repeatedly isolated (60 of 140) detected transcripts only in RNA prepared from Th2 clones (D10, CDC35) and not from either Th1 clones (AR5, OS6, D1) or from a B cell lymphoma, M12, as illustrated in the Northern blot analysis depicted in Figure 2A. Further, the levels of transcripts detected in D10 Th2 cells were substantially increased upon activation by ligation of the T cell receptor with anti-CD3 antibody. No induction of the transcript detected by this cDNA clone occurred in Th1 clones upon anti-CD3 treatment. A control probe, GAPDH, demonstrated approximately equal loading of RNA in all lanes. Thus, the expression of this cDNA clone in the lymphoid lineage appeared to be Th2-specific and sensitive to signals transmitted through the T cell receptor. For these Northern blots, total RNA was prepared by using Trizol (GIBCO/BRL) according to manufacturer's instructions. 10 µg of total RNA from each sample was fractionated on a formaldehyde agarose gel and transferred to a nylon membrane. A 300 bp *Dra*I fragment derived from the 3' untranslated region of the isolated clone was labeled with  $\alpha$ -<sup>32</sup>P-dCTP using Random Primed DNA Labeling Kit (Boehringer Mannheim, Indianapolis, IN). Hybridization was performed using QuikHyb (Stratagene, La Jolla, CA) according to manufacturer's instructions.

To determine whether the expression of this gene was tissue-specific and regulated during the course of normal Th cell development, the following experiment was performed. Naive spleen cells (Th precursor (Thp) cells) were driven along a Th1 or Th2 pathway by

66248756 "03499  
treatment with anti-CD3 in the presence of cytokines and anti-cytokine antibodies (IFN $\gamma$  and anti-IL-4 for Th1, IL-4 and anti-IFN $\gamma$  for Th2). Splenic cell suspensions were prepared from 6-8 week-old Balb/c mice, cultured in RPMI 1640 supplemented with 10% FCS at a density of 10<sup>6</sup> cells/ml, and stimulated with plate bound anti-CD3 antibody in the presence of  
5 5  $\mu$ g/ml of anti-IL4 antibody (11B11) for the Th1 lineage, or 5  $\mu$ g/ml of anti-IFN $\gamma$  antibody (XMG-1) for the Th2 lineage. 24 hours after stimulation, 50 U/ml IL2 was added to all cultures, and 500 U/ml IL4 (Genzyme) was added to Th2 cultures. 7 days after the primary stimulation, all cells were harvested, washed and restimulated with plate bound anti-CD3 antibody. Northern blot analysis of differentiating cells harvested at various time points after  
10 stimulation in a primary (day 0-8) and secondary (0-20 hours) response was performed, using the methodology described above, and identification of differentiating Thp cells as Th1 or Th2 was determined by analyzing culture supernatants by ELISA for IL-10 and IFN $\gamma$ . ELISA for cytokine quantitation was performed as follows. All anti-cytokine antibodies were purchased from Pharmingen. ELISA was performed according to Pharmingen's  
15 instructions with the exception that Avidin-Alkaline Phosphatase (Sigma) at 1:500 dilution in PBS/BSA was used in place of avidin-peroxidase. P-nitrophenyl phosphate (GIBCO BRL) at 4 mg/ml in substrate buffer (10% diethanolamine, 0.5 mM MgCl<sub>2</sub>, 0.02% sodium azide, pH 9.8) was used as substrate.

In two independent experiments, representative results of which are shown in Figure  
20 2B, this analysis revealed low level or undetectable expression of this cDNA in naive spleen cells at baseline at day 0. In cultures differentiating along a Th2 pathway, substantial induction of transcripts occurred by day 8 in a primary stimulation and by 20 hr in a secondary stimulation. In contrast, no induction occurred in cells being driven along a Th1 pathway. A control probe (GAPDH) showed approximately equal loading of RNA in all  
25 lanes. The low level of transcripts present in cells being driven along a Th1 pathway likely reflects the presence of residual Th2 cells since complete skewing does not occur in this *in vitro* differentiation system.

Together, these experiments revealed that the isolated cDNA is selectively expressed in Th2 clones, where it is induced upon T cell activation, and that it is absent from Th1  
30 clones and a B lymphoma. Further, this gene is induced in normal Thp when they are driven towards the Th2 lineage, but is not induced during Th1 development.

The cDNA obtained from the yeast two-hybrid screen was used as a probe to isolate a full-length cDNA from a D10 Th2 cell cDNA library by standard hybridization methods. A 4.3 kb cDNA clone was isolated from the Th2 cell library and sequenced by standard  
35 methods. Sequence analysis revealed that this Th2-specific gene corresponded in sequence to the *c-maf* proto-oncogene.

**EXAMPLE 3: Ectopic Expression of c-Maf in Th1 and B Cells Results in Activation of the IL-4 Promoter**

The identification of the isolated cDNA described in Example 2 as a member of the AP-1/CREB/ATF gene family, together with its selective expression in Th2 cells raised the possibility that c-Maf controlled the tissue-specific transcription of the IL-4 gene. Additionally, the presence of transcripts encoding c-maf correlated well with IL-4 expression in Th2 cells and in three of four transformed mast cell lines examined. To test whether c-Maf could transactivate the IL-4 promoter, cotransfection experiments were performed.

Th1 clones and the B lymphoma M12.4.C3 (M12) neither express *c-maf* nor transcribe the IL-4 gene. If c-Maf is the transcription factor critical for controlling IL-4 gene expression, then forced expression in these cells should permit IL-4 gene expression. To test this, the full-length (4.3 kb) c-maf cDNA clone was inserted into the SalI site of the pMex-Neol mammalian expression vector, which utilizes the CMV enhancer to drive expression of the inserted sequence. The c-Maf expression vector was then cotransfected with an IL-4 promoter reporter construct into the Th1 clone AE7 and the B lymphoma M12. The generation of the wild type IL4 CAT reporter construct, containing an IL4 promoter fragment from -157 to +68 operatively linked to a chloramphenicol acetyltransferase gene is described in Hodge, M. *et al.* (1995) *J. Immunol.* 154:6397-6405. The Th1 clone was cultured in RPMI 1640 supplemented with 10% FCS and 10% Con-A stimulated rat splenocyte supernatant, and maintained by bi-weekly stimulation with appropriate antigen and APCs. M12 cells were cultured in RPMI 1640 supplemented with 10% FCS.

The Th1 clone AE7 or M12 B lymphoma cells were transiently transfected by preincubating 0.4 ml of cells, containing  $2 \times 10^7$  cells/ml AE7 or  $3 \times 10^6$  cells/ml M12 cells in serum-free RPMI 1640 with 20  $\mu$ g (AE7) or 5  $\mu$ g (M12) of each plasmid for 10 minutes at room temperature. The samples were then electroporated using a BIO-RAD Gene Pulser (BIO-RAD, Richmond, CA) set at 975  $\mu$ F, 280 V, and immediately placed on ice for 10 minutes. The transfected cells were allowed to recover overnight in complete media and stimulated with plate bound anti-CD3 antibody ({Pharmingen, San Diego, CA} 1  $\mu$ g/ml in 1XPBS overnight at 4 °C) or with 50 ng/ml PMA (Sigma, St. Louis, MO) and 1  $\mu$ M Ionomycin (Calbiochem Corp., La Jolla, California). for 24 hours. Cell lysate was prepared by freeze-thaw lysis in 0.25 M Tris-Cl, pH 7.8. Equal amounts of protein (between 5-20  $\mu$ g) were used for CAT assays. CAT assays were performed as described in Todd, M. *et al.* (1993) *J. Exp. Med.* 177:1663-1674.

It has previously shown that Th2-specific, inducible IL-4 expression can be directed by as little as 157 bp of the proximal IL-4 promoter in Th2 cells (Hodge, M. *et al.* (1995) *J. Immunol.* 154:6397-6405). In cotransfection experiments, the results of which are summarized in Figure 3A, it is demonstrated that ectopic expression of c-Maf in the Th1 clone AE7 results in substantial activity of the IL-4 promoter reporter after stimulation

through the T cell receptor. The fold induction observed was approximately 5 fold over that observed with the control empty vector alone. Although expression of a reporter construct containing proximal (-157 to +58) IL-4 promoter sequences in the subclone of AE7 cells utilized here has not been previously observed, it has been demonstrated that small amounts of IL-4 mRNA can be detected by RT-PCR in other subclones of AE7. To more rigorously test the ability of c-Maf to transactivate the IL-4 promoter in a non-IL-4 producing cell, the same experiment was performed in the B lymphoma cell line, M12. Normal B cells and B lymphoma cells do not produce IL-4. Representative results of the cotransfection experiments are depicted in Figure 3B and a summary of three independent experiments is shown below in Table 1.

**Table 1**

Plasmids	CAT Activity (fold induction)			
	PMA/iono.	Exp. I*	Exp. II	Exp. III
pMEX-NeoI/pREP4	-	1	1	1
	+	7.6	1	1.4
pMEX-Maf/pREP4	-	95	5	18.6
	+	186	7	37
pMEX-c-Fos/pREP4	-	2.7	1	0.8
	+	7.6	1.2	1
pMEX-JunD/pREP4	-	ND**	0.9	0.5
	+	ND	1.4	1.9
pMEX-NeoI/pREP4-NFATp	-	14.2	1.6	0.3
	+	41.2	3.5	0.3
pMEX-Maf/pREP4-NFATp	-	136	54	26.3
	+	138	100	54.7
pMEX-c-Fos/pREP4-NFATp	-	7.4	1.6	3
	+	15.4	1.9	6.1

\* In experiment I, 20 mg of cell lysate was incubated for 2 hours. In experiments II and III, only 5 mg of cell lysate was incubated for 1 hour in order to reveal synergy between c-Maf and NFATp

\*\*ND = not done

The results in M12 B lymphoma cells confirmed the findings in the Th1 clone. Ectopic expression of c-Maf resulted in substantial activity of the IL-4 promoter in M12 cells, either unstimulated or stimulated with PMA/Ca<sup>++</sup> ionophore. The fold induction observed when compared to transfection of a control vector averaged approximately 50 in unstimulated M12 cells. Stimulation of M12 cells with PMA/Ca<sup>++</sup> ionophore, which should result in translocation of NF-ATs to the nucleus and induction of other AP-1 family members (Flanagan, W.M. (1991) *Nature* 352:803-807; Jain, J. *et al.* (1993) *Nature* 365:352-355), increased the basal activity of the IL-4 promoter, but a marked induction in promoter activity

by c-Maf was still present (average of approximately 25 fold). C-Maf did not transactivate a control reporter driven by NF-AT multimers, demonstrating the specificity of transactivation.

As a control for the specificity of c-Maf as opposed to other AP-1 family members, the c-Fos and c-Jun proteins were also overexpressed in M12 cells utilizing murine full-length cDNAs encoding c-Fos and JunD in the mammalian expression vector of pMEX-NeoI together with the IL-4 reporter plasmid. No IL-4 promoter activity could be achieved by overexpression of either of these two AP-1 family members in M12 cells. Thus, c-Maf has a unique ability to drive IL-4 gene transcription in M12 B cells. Further, forced expression of c-Maf in the hepatoma cell line HepG2 also resulted in IL-4 promoter transactivation. These experiments demonstrate that the provision of c-Maf to c-Maf negative Th1 or B cells, or to non-lymphoid cells (*e.g.*, a hepatoma cell line), permits the cells to transactivate the IL-4 promoter.

NF-AT proteins have been shown to be critically important in the regulation of both the IL-4 and IL-2 cytokines. NF-ATp was the first member of this family to be isolated (McCaffrey, P.G. *et al.* (1993) *Science* 262:750-754). Both AE7 and M12 cells have endogenous NF-ATp protein, but nevertheless do not transcribe IL-4. Although NF-ATp could not therefore account for selective IL-4 gene transcription, it was of interest to test whether overexpression of NF-ATp in unstimulated or stimulated M12 cells would further increase the transactivation of the IL-4 promoter by c-maf. M12 cells were cotransfected with the IL-4 reporter construct and either an NFAPp expression vector (pREP4-NF-ATp, which also carries a hygromycin resistance gene) alone or the NFAPp expression vector together with the c-Maf expression vector. Overexpression of NF-ATp alone in M12 cells resulted in some modest transactivation of the IL-4 promoter. This transactivation was markedly increased by ectopic expression of c-Maf, an increase which was not just additive but was synergistic (see Fig. 3B and Table 1). In contrast, c-Fos overexpression did not further increase the modest transactivation achieved by NF-ATp. These results indicate that c-maf and NF-ATp interact to achieve maximal induction of the IL-4 promoter, the tissue-specificity being provided by c-maf.

**EXAMPLE 4: Ectopic Expression of C-Maf Activates Transcription of the Endogenous IL-4 Gene in a B Lymphoma**

As demonstrated in Example 3, c-Maf transactivates the IL-4 promoter in transient transfection assays in Th1, B and non-lymphoid cells. To test whether expression of c-maf in non-IL-4 producing cells can activate the transcription of endogenous IL-4, the B lymphoma M12 was stably transfected with expression vectors encoding c-maf, NF-ATp or both, or junD with and without NF-ATp as a control. For stable transfection, M12 cells were transfected as described above in Example 3. The transfected cells were allowed to recover in complete media for 48 hours before the addition of Neomycin (GIBCO/BRL,

Gaithersburg, MD) and Hygromycin (Calbiochem, Corp.) at a concentration of 400 µg/ml of each antibiotic. The transfected cells were supplemented with fresh media every other day.

Stably transfected M12 cells were plated at equal density supernatants harvested 24 hours later to measure cytokines by ELISA. ELISAs were performed as described in Example 2. The results, shown in Figure 4, demonstrate that in these experiments M12 cells transfected with c-maf, junD or NF-ATp alone did not produce measurable IL-4 by ELISA. However, M12 cells stably transfected with both c-maf and NF-ATp did produce detectable, but low level, IL-4 by ELISA. These results were confirmed by RT-PCR on RNA from these transfected cells. In contrast, these cells did not produce detectable IL-2. The requirement for both c-maf and NF-ATp is consistent with the synergistic effect of these factors in the transactivation of the IL-4 promoter noted in the transient transfection experiments in M12 cells. In contrast, transfection of junD, an AP-1 family member which can increase IL-4 expression in Th2 cells, alone or together with NF-ATp, did not result in IL-4 production. These results demonstrate the essential and selective role of c-maf in directing tissue-specific endogenous IL-4 production.

**EXAMPLE 5:        A Site in the IL-4 Promoter is Footprinted by  
                         Extracts from Th2 but not Th1 Clones**

The experiments described in Examples 3 and 4 demonstrated a clear functional role for c-maf in controlling tissue-specific expression of IL-4. Further, c-maf transcripts were expressed in Th2 but not Th1 cells. However, DNA-protein complexes were not detected by electrophoretic mobility shift assays (EMSA) when using nuclear extracts prepared from Th2 cells. To further examine whether a protein in Th2 nuclear extracts might bind to the MARE, or nearby sequences, the more sensitive technique of DNaseI footprinting was used. Two Th2 clones (D10, CDC35) and two Th1 clones (AE7, S53) were activated by ligation of the T cell receptor with plate-bound anti-CD3 antibody, and nuclear extracts prepared at time 0 (unstimulated), 2 hours and 6 hours later. DNaseI footprinting analysis was then performed according to standard methods using a Klenow end-labeled IL-4 promoter fragment (-157 to +68). The results are shown in Figure 5A. Stimulated extracts from both Th1 and Th2 cells footprinted the two NF-AT sites and the AP-1 site upstream of the distal NF-AT site as described previously (Rooney, J. *et al.* (1995) *Immunity* 2:545-553), consistent with the demonstrated function of NF-AT and AP-1 proteins in regulating both the IL-2 and the IL-4 promoters (Rooney, J. *et al.* (1995) *Immunity* 2:545-553; Rooney, J. *et al.* (1995) *Mol. Cell Biol.* 15:6299-6310). Furthermore, inspection of the autoradiograph revealed an area of hypersensitivity on the non-coding strand at residues -28 and -29 when extracts from stimulated Th2 but not stimulated Th1 cells were used. Unstimulated Th cell extracts did not footprint this region. The Th2 footprint observed was subtle, but reproducible in two experiments and is located in a site that has previously been demonstrated to be critical for

IL-4 promoter activation in Th2 cells (Hodge, M. *et al.* (1995) *J. Immunol.* 154:6397-6405). A schematic summary of sites occupied in the IL-4 promoter as detected by footprint analysis is shown in Figure 5B. These results indicate that a site in the proximal IL-4 promoter, previously shown to be functionally important, is occupied in activated Th2 but not in activated Th1 cells.

**EXAMPLE 6: Recombinant c-maf Binds to a MARE Site in the IL-4 Promoter**

The Th2-specific footprint does not contain a c-maf response element (MARE). However, examination of the proximal IL-4 promoter revealed a half c-maf binding site (MARE) ( residues -42 to -37) immediately downstream of the proximal NF-AT site (residues -56 to -51) (shown schematically in Fig 5B). It has previously been demonstrated that mutation of this site abolished activity of the IL-4 promoter in Th2 cells (Hodge, M. *et al.* (1995) *J. Immunol.* 154:6397-6405). To determine if c-Maf bound this site, a truncated c-Maf recombinant protein containing the b-zip domain(amino acids 171-371) was expressed from *E. coli*, purified on an S-Tag agarose column and used in electrophoretic mobility shift assays with radiolabeled MARE oligonucleotide.

The expression vector for recombinant c-Maf was constructed by inserting a cDNA fragment encoding a.a. residues 171 to 371 of c-Maf (disclosed in Kurschner C. and Morgan, J.I. (1995) *Mol. Cell. Biol.* 15:246-254) into the NotI site of pET29 (Novagen, Inc. Madison, WI). The truncated c-Maf protein was expressed using T7 polymerase in the BL21(DE3)strain. Cells were induced by the addition of 1 mM IPTG and incubated at 37° C for 3 hours. The induced cells were lysed in 1 X Bind/Wash buffer (20 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% Triton X-100) followed by sonication. The c-Maf protein was then purified from the soluble fraction by using the S-Tag Purification Kit (Novagen) according to manufacturer's instructions. Two additional proteins, NF-ATp and c-Jun, were also used in EMSA assays. The recombinant NF-ATp, containing the Rel domain of murine NF-ATp, was expressed using an *in vitro* transcription/translation vector TP7-NF-ATp, which contains a cDNA fragment encoding the Rel domain of murine NF-ATp. The c-Jun expression vector, pGEM-c-Jun, was constructed by inserting a full-length cDNA of murine c-Jun into the PstI site of pGEM4. 1 µg of each plasmid DNA was transcribed from the T7 promoter and translated in rabbit reticulocyte lysate by using the TnT Coupled Transcription/Translation Kit (Promega, Madison, WI).

Electrophoretic mobility shift assays (EMSA) were performed as follows. 100 ng of double-stranded oligonucleotides were end-labeled with  $\gamma$ -<sup>32</sup>P-dATP (DuPont NEN Research Product, Wilmington, DE) using T4 polynucleotide kinase (Pharmacia LKB Biotechnology, Inc., Piscataway, NJ). The labeled ds-oligonucleotides were fractionated on 15-20% polyacrylamide gels, eluted overnight at 37° C in 1X TE and precipitated in ethanol. Binding assays were performed at room temperature for 20 minutes using 0.5 µg of recombinant



proteins or 4 µl of *in vitro* translated products, 500 ng poly(dI-dC), and 20,000 cpm of probe in a 15 µl volume of 20 mM HEPES (pH 7.9), 100 mM KCl, 5% glycerol, 1mM EDTA, 5mM DTT, 0.1% NP-40, and 0.5 mg/ml BSA. The samples were then fractionated in 4% non-denaturing polyacrylamide gel containing 0.5X TBE at room temperature.

5 Oligonucleotides derived from the murine IL4 promoter used in EMSA were:

-59 to -27: 5'-CTCATTTTCCCTTGGTTTCAGCAACTTTAACTC-3' (SEQ ID NO: 1);

-79 to -60: 5'-ATAAAATTTTCCAATGTAAA-3' (SEQ ID NO: 2); and

-88 to -61: 5'-TGGTGTAATAAAATTTTCCAATGTAAA-3' (SEQ ID NO: 3).

The sequence of the MARE oligonucleotide used in EMSA was:

10 5'-GGAATTGCTGACTCAGCATTACT-3'(SEQ ID NO: 4).

All oligonucleotides were annealed with their respective reverse-complementary strands to form double-stranded oligonucleotides.

The results of EMSA with recombinant c-Maf are shown in Figure 6. The recombinant c-Maf protein bound well to both a consensus MARE oligonucleotide and to a  
15 33 bp oligonucleotide containing the NF-AT site and MARE present in the IL-4 promoter. Binding was specifically competed by unlabeled homologous but not control probe. Further, c-Maf did not bind to an oligonucleotide containing only the NF-AT target sequence to which recombinant NF-ATp bound well. The ability of c-Maf to bind to the IL-4 promoter probe was specific since *in vitro* translated c-Jun protein did not bind to this oligonucleotide. The  
20 c-Jun protein was functional since it could bind to the consensus MARE which contains a core TRE site. These results indicate that c-Maf, but not another AP-1 family member (c-Jun), can bind to the MARE site within the proximal IL-4 promoter.

NF-AT proteins interact cooperatively with AP-1 family member proteins to form higher mobility complexes on IL-2 and IL-4 promoter DNA on EMSA (Jain, J. (1993)  
25 *Nature* 365:353-355; Rooney, J. *et al.* (1995) *Immunity* 2:545-553). That NF-AT proteins might interact with c-maf was suggested by the functional studies described in the previous examples. To determine if c-Maf interacted with NF-AT in the presence of DNA, recombinant NF-ATp and c-Maf proteins were used separately or together in EMSA with the  
30 33 bp oligonucleotide containing both the NF-AT and adjacent MARE sites. The results are shown in Figure 6. Each protein alone bound to IL-4 promoter DNA. Recombinant c-Maf plus recombinant NF-ATp protein produced these complexes and in addition formed a higher mobility complex. No higher mobility complex was observed when c-Jun and NF-ATp proteins were used, consistent with the failure of c-Jun to bind this site. These results indicate that c-Maf can specifically bind *in vitro* to a sequence located in the proximal IL-4  
35 promoter, previously shown to be functionally critical in Th2 cells, and that, like other AP-1 proteins, c-Maf can interact *in vitro* with NF-AT proteins.

**EXAMPLE 7:            The Ability of c-Maf to Transactivate the IL-4 Promoter**  
**Maps to the MARE and Th-2 Specific Footprint**

5        An essential region of the IL-4 promoter located immediately upstream of the TATA  
element has been characterized by high resolution mutagenesis (Hodge, M. *et al.* (1995) *J.*  
*Immunol.* 154:6397-6405). Mutagenesis of this 33 bp region (-59 to -28) demonstrated  
multiple sites required for inducible IL-4 transcription in Th2 cells. These sites included an  
NF-AT target sequence, the region footprinted by Th2 extracts, and what is now recognized  
as a MARE. A series of IL-4 reporter gene constructs comprising 4 base pair linker-scanning  
10       mutants generated across this region were used to map the target sequence utilized by c-Maf  
*in vivo* in M12 cells. These cells were cotransfected with the c-maf expression vector and  
this series of mutant IL-4 promoter constructs. The results are shown in Figure 7A.  
Mutation of the MARE (mut 3 and 4), or the site defined by the Th2 footprint (mut 2),  
abrogated (mut 2 and 4) or partially abrogated (mut 3) the ability of transfected c-maf to  
15       drive IL-4 transcription. A modest effect in reducing c-maf transactivation was also observed  
for mutant 8 which disrupts the NF-AT sequence, consistent with the presence in M12 cells  
of endogenous NF-ATp and with the synergy between NF-ATp and c-maf demonstrated in  
the previous examples. Mutants 6 and 7 had no significant effect while mutant 5 had  
enhanced transactivation ability, consistent with previous observations in Th2 cells (Hodge,  
20       M. *et al.* (1995) *J. Immunol.* 154:6397-6405). The transactivation data is consistent with  
EMSA performed with recombinant c-Maf protein using as probe an oligonucleotide which  
contains this 33 bp region, and this same series of mutant oligonucleotides as cold  
competitors. The results of these EMSA experiments are shown in Figure 7B. These  
experiments indicate that c-Maf specifically binds to and transactivates the MARE in the  
25       proximal IL-4 promoter and that the adjacent Th2-specific element is intimately involved in  
both the binding and function of c-Maf.

**EQUIVALENTS**

30       Those skilled in the art will recognize, or be able to ascertain using no more than  
routine experimentation, many equivalents to the specific embodiments of the invention  
described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Glimcher, Laurie H.  
Ho, I-Cheng

10

(ii) TITLE OF INVENTION: Methods and Compositions for Regulating T Cell  
Subsets by Modulating Transcription Factor  
Activity

(iii) NUMBER OF SEQUENCES: 4

15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: LAHIVE & COCKFIELD  
(B) STREET: 60 State Street, suite 510  
(C) CITY: Boston  
(D) STATE: Massachusetts  
(E) COUNTRY: USA  
(F) ZIP: 02109-1875

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US  
(B) FILING DATE:  
(C) CLASSIFICATION:

30

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: DeConti, Giulio A., Jr.  
(B) REGISTRATION NUMBER: 31,503  
(C) REFERENCE/DOCKET NUMBER: HUI-021

35

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617)227-7400  
(B) TELEFAX: (617)227-5941

40

(2) INFORMATION FOR SEQ ID NO:1:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

55

CTCATTTTCC CTTGGTTTCA GCAACTTTAA CTC

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATAAAATTTT CCAATGTAAA

20

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGGTGTAATA AAATTTTCCA ATGTAAA

27

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGAATTGCTG ACTCAGCATT ACT

23

662130 "95284260

CLAIMS

1. A method for modulating production of a T helper type 2 (Th2)-associated cytokine by a cell comprising contacting the cell with an agent that modulates the expression or activity of a transcription factor that regulates expression of a Th2-associated cytokine gene such that production of the Th2-associated cytokine by a cell is modulated.

2. The method of claim 1, wherein the agent acts intracellularly to modulate the expression or activity of the transcription factor that regulates expression of a Th2-associated cytokine gene.

3. The method of claim 1, wherein the transcription factor is a member of the maf family.

4. The method of claim 3, wherein the transcription factor is c-Maf.

5. The method of claim 1, wherein the Th2-associated cytokine is interleukin-4.

6. The method of claim 1, further comprising contacting the cell with a second agent that modulates the expression or activity of a second transcription factor that contributes to regulating the expression of a Th1- or Th2-associated cytokine gene.

7. The method of claim 6, wherein the second agent modulates the expression or activity of a Nuclear Factor of Activated T cells.

8. The method of claim 1, wherein production of a Th2-associated cytokine by the cell is stimulated.

9. The method of claim 8, wherein the cell is a T helper type 1 (Th1) cell, a B cell or a nonlymphoid cell.

10. The method of claim 8, wherein the agent is a nucleic acid molecule encoding a maf family protein, wherein the nucleic acid molecule is introduced into the cell in a form suitable for expression of the maf family protein in the cell.

11. The method of claim 1, wherein production of a Th2-associated cytokine by the cell is inhibited.

12. The method of claim 11, wherein the cell is a Th2 cell.

13. The method of claim 11, wherein the agent is an intracellular binding molecule.

5 14. The method of claim 1, further comprising administering the cell to a subject to thereby modulate development of T helper type 1 (Th1) or T helper type 2 (Th2) cells in a subject.

10 15. A method for modulating development of T helper type 1 (Th1) or T helper type 2 (Th2) cells in a subject comprising administering to the subject an agent that modulates the activity of a transcription factor that regulates expression of a Th2-associated cytokine gene such that development of Th1 or Th2 cells in the subject is modulated.

15 16. The method of claim 15, wherein the agent acts intracellularly to modulate the expression or activity of the transcription factor that regulates expression of a Th2-associated cytokine gene.

20 17. The method of claim 15, wherein the transcription factor is a member of the maf family.

18. The method of claim 17, wherein the transcription factor is c-Maf.

19. The method of claim 15, wherein the Th2-associated cytokine is interleukin-4.

25 20. The method of claim 15, further comprising administering to the subject a second agent that modulates the expression or activity of a second transcription factor that contributes to regulating the expression of a Th1- or Th2-associated cytokine gene.

30 21. The method of claim 20, wherein the second agent modulates the expression or activity of a Nuclear Factor of Activated T cells.

22. The method of claim 15, wherein production of a Th2-associated cytokine by cells of the subject is stimulated.

35 23. The method of claim 15, wherein production of a Th2-associated cytokine by cells of the subject is inhibited.

24. A recombinant expression vector comprising a nucleotide sequence encoding a maf family protein operatively linked to regulatory sequences that direct expression of the maf family protein specifically in lymphoid cells.

5 25. The recombinant expression vector of claim 24, wherein the regulatory sequences direct expression of the maf family protein specifically in T cells.

26. The recombinant expression vector of claim 24, wherein the regulatory sequences direct expression of the maf family protein specifically in B cells.

10

27. A recombinant expression vector comprising a nucleotide sequence encoding a maf family protein operatively linked to regulatory sequences that direct expression of the maf family protein specifically in hematopoietic stem cells.

15 28. A host cell into which a recombinant expression vector encoding a maf family protein has been introduced, wherein the host cell is a lymphoid cell.

29. The host cell of claim 28, which is a T cell.

20 30. The host cell of claim 28, which is a B cell.

31. A host cell into which a recombinant expression vector encoding a maf family protein has been introduced, wherein the host cell is a hematopoietic stem cell.

25 32. A method for identifying a compound that modulates the expression or activity of a transcription factor that regulates expression of a Th2-associated cytokine gene comprising:

- 30 a) preparing an indicator cell, wherein said indicator cell contains:  
i) a recombinant expression vector encoding a transcription factor that regulates expression of a Th2-associated cytokine gene; and  
ii) a vector comprising regulatory sequences of the Th2-associated cytokine gene operatively linked a reporter gene;
- 35 b) contacting the indicator cell with a test compound;  
c) determining the level of expression of the reporter gene in the indicator cell in the presence of the test compound;
- d) comparing the level of expression of the reporter gene in the indicator cell in the presence of the test compound with the level of expression of the reporter gene in the indicator cell in the absence of the test compound; and

e) identifying a compound that modulates the expression or activity of a transcription factor that regulates expression of a Th2-associated cytokine gene.

33. A method for identifying a protein in a Th2 cell that interacts with c-Maf  
5 comprising:
- a) providing a two hybrid assay including a host cell which contains
    - i) a reporter gene operably linked to a transcriptional regulatory sequence;
    - ii) a first chimeric gene which encodes a first fusion protein, said first  
10 fusion protein including c-Maf;
    - iii) a library of second chimeric genes which encodes second fusion proteins, the second fusion proteins including proteins derived from Th2 cells;
- wherein expression of the reporter gene is sensitive to interactions between the first  
15 fusion protein, the second fusion protein and the transcriptional regulatory sequence;
- b) determining the level of expression of the reporter gene in the host cell;
- and
- c) identifying a protein in a Th2 cell that interacts with c-Maf.

- 20 34. A method for identifying a compound that modulates the interaction of a c-Maf protein with a maf response element (MARE) of an IL-4 gene regulatory region, comprising:
- a) providing a c-Maf protein and a DNA fragment comprising a MARE of an IL-4 gene regulatory region;
  - 25 b) incubating the c-Maf protein and DNA fragment in the presence of a test compound;
  - c) determining the amount of binding of the c-Maf protein to the DNA fragment in the presence of the test compound;
  - d) comparing the amount of binding of the c-Maf protein to the DNA fragment in the  
30 presence of the test compound with the amount of binding of the c-Maf protein to the DNA fragment in the absence of the test compound; and
  - e) identifying a compound that modulates the interaction of a c-Maf protein with a MARE of an IL-4 gene regulatory region.



**METHODS AND COMPOSITIONS FOR REGULATING T CELL SUBSETS BY  
MODULATING TRANSCRIPTION FACTOR ACTIVITY**

5

**Abstract**

Methods for modulating production of a T helper type 2 (Th2)-associated cytokine, in particular interleukin-4, by modulating the activity of a transcription factor, in particular the proto-oncoprotein c-Maf, that regulates expression of the Th2-associated cytokine gene are disclosed. Methods for modulating development of T helper type 1 (Th1) or T helper type 2 (Th2) subsets in a subject using agents that modulate transcription factor activity are also disclosed. The methods of the invention can further involve use of agents that modulate the activity of additional transcription factors that contribute to the regulation of Th1- or Th2-associated cytokines, such as a Nuclear Factor of Activated T cells (NF-AT) protein and/or an AP-1 family protein. Compositions for modulating Th2-associated cytokine production, recombinant expression vectors and host cells, as well as screening assays to identify agents that modulate c-Maf activity, are also disclosed.

0949795284260




FIGURE 1A




FIGURE 1B




FIGURE 2A




FIGURE 2B




FIGURE 3A

662120" 95284250




FIGURE 3B




FIGURE 4






FIGURE 5A

-100                      -80                      -60                      -40                      -20  
 GAATAACTGACAATCTG**GTGTAATAAA**TTTTC**CAATGTA**AACTCATT**TTTCCCTTGGTTTCAGCA**ACTTTAACTCTATATATA  
 CTTATTGACTGTTAGACCACATTATTT**TAAAAGG**TTACATTGAGT**TAAAAGGGAACCA**AGTCGTTGAAATTGAGATATATAT  
                                  AP-1                      NFAT                      NFAT                      MARE                      \*\*  
    Th2-specific  
    footprint

WT -27 to -59  
 mut 2  
 mut 3  
 mut 4  
 mut 5  
 mut 6  
 mut 7  
 mut 8

CTCATTTTCCCTTGGTTTCAGCAACTTTAACTC  
 CTCATTTTCCCTTGGTTTCAGCAAC**gggc**ACTC  
 CTCATTTTCCCTTGGTTTCAG**acca**TTTAACTC  
 CTCATTTTCCCTTGGTT**gact**CAACTTTAACTC  
 CTCATTTTCCCTT**ttgg**TCAGCAACTTTAACTC  
 CTCATTTT**Caagg**GGTTTCAGCAACTTTAACTC  
 CTCAT**ggga**CCTTGGTTTCAGCAACTTTAACTC  
 C**gacg**TTTCCCTTGGTTTCAGCAACTTTAACTC

FIGURE 5B




FIGURE 6




FIGURE 7A




FIGURE 7B

Attorney's  
Docket  
Number HUI-021

Declaration, Petition and Power of Attorney For Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**METHODS AND COMPOSITIONS FOR REGULATING T CELL SUBSETS BY  
MODULATING TRANSCRIPTION FACTOR ACTIVITY**

the specification of which

(check one)

is attached hereto.

was filed on April 23, 1996 as

Application Serial No. 08/636,602

and was amended on \_\_\_\_\_  
(if applicable)

I do not know and do not believe that the subject matter of this application was ever known or used in the United States before my invention thereof or patented or described in any printed publication in any country before my invention thereof or more than one year prior to the date of this application, and that said subject matter has not been patented or made the subject of an issued inventor's certificate in any country foreign to the United States on an application filed by me or my legal representatives or assigns more than twelve months prior to the date of this application; that I acknowledge my duty to disclose information of which I am aware which is material to the examination of this application, that no application for patent or inventor's certificate on the subject matter of this application has been filed by me or my representatives or assigns in any country foreign to the United States, except those identified below, and that I have reviewed and understand the contents of the specification, including the claims as amended by any amendment referred to herein.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

09248756-031299  
66729-92420

## PRIORITY CLAIM

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed.

Check one:

☒ no such applications have been filed.

☐ such applications have been filed as follows

EARLIEST FOREIGN APPLICATION(S), IF ANY FILED WITHIN 12 MONTHS  
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

Country	Application Number	Date of Filing (month,day,year)	Priority Claimed Under 35 USC 119
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No

ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTHS  
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION


# CLAIM FOR BENEFIT OF EARLIER U.S./PCT APPLICATION(S)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application. As to subject matter of this application which is common to my earlier United States application, if any, described below, I do not believe that the same was ever known or used in the United States before my invention thereof or patented or described in any printed publication in any country before my invention thereof or more than one year prior to said earlier application, or in public use or on sale in the United States more than one year prior to said earlier application, that the said common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of said earlier application in any country foreign to the United States on an application, filed by me or my legal representatives or assigns more than twelve months prior to said application and that no application for patent or inventor's certificate on said subject matter has been filed by me or my representatives or assigns in any country foreign to the United States except those identified herein.

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented,pending,aband.)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented,pending,aband.)



POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

John A. Lahive, Jr.	Reg. No. 19,788	John V. Bianco	Reg. No. 36,748
W. Hugo Liepmann	Reg. No. 20,407	Jeremiah Lynch	Reg. No. 17,425
James E. Cockfield	Reg. No. 19,162	Amy E. Mandragouras	Reg. No. 36,207
Thomas V. Smurzynski	Reg. No. 24,798	Elizabeth A. Hanley	Reg. No. 33,505
Ralph A. Loren	Reg. No. 29,325	Anthony A. Laurentano	Reg. No. 38,220
Thomas J. Engellenner	Reg. No. 28,711	Jane E. Remillard	Reg. No. 38,872
Giulio A. DeConti, Jr.	Reg. No. 31,503	Mark A. Kurisko	Reg. No. 38,944
Ann Lamport Hammitte	Reg. No. 34,858	Beth E. Arnold	Reg. No. 35,430
Paul Louis Myers	Reg. No. 35,965	Jean M. Silveri	Reg. No. 39,030
Michael I. Falkoff	Reg. No. 30,833	Matthew P. Vincent	Reg. No. 36,709

Send Correspondence to:


Lahive & Cockfield, 60 State Street, Suite 510, Boston, MA 02109-1875


Direct Telephone Calls to: (name and telephone number)

Giulio A. DeConti, Jr., Esq., (617) 227-7400

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor	Laurie H. Glimcher	
Inventor's signature		Date 6/3/96
Residence	51 Hampshire Street, West Newton, Massachusetts 02165	
Citizenship	U.S.A.	
Post Office Address (if different)		

Full name of second inventor, if any I-Cheng Ho	
Inventor's signature 	Date 6/5/96
Residence 30 Silver Birch Road, Newton, Massachusetts 02168	
Citizenship Taiwan	
Post Office Address (if different)	

09349266-03202020

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the application of: Laurie H. Glimcher et al.

Serial No.: Continuation of 08/636,602

Filed: herewith

For: Methods and Compositions For Regulating T  
Cell Subsets By Modulating Transcription Factor  
Activity

Attorney Docket No.: HUI-021CN

Group Art Unit: 1632

Examiner: S. Priebe

Assistant Commissioner for Patents  
Washington, D.C. 20231

**ASSOCIATE POWER OF ATTORNEY**


Sir:

The undersigned attorney has the power of attorney in the subject application. He  
hereby grants an associate power to:

Megan E. Williams, Ph.D.  
Registration No. 43,270  
Lahive & Cockfield, LLP  
28 State Street  
Boston, MA 02109

Please continue to forward all written and telephonic communications to Amy E.  
Mandragouras at the address and telephone number listed below.

Respectfully submitted,

  
Giulio A. DeConti, Jr.  
Registration No. 31,503  
Attorney for Applicants

LAHIVE & COCKFIELD, LLP  
28 State Street  
Boston, MA 02109  
Tel. (617) 227-7400  
Dated: February 12, 1999